


CSIR-UGC-NET Chemical Science - June - 2018



- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching
- ✓ PG | Polytechnic TRB Chemistry Coaching

www.csircoaching.com

Features


- | | |
|-------------------------------|----------------------------------|
| ➤ 300 ++ Live Classes | ➤ A Well-Defined Curriculum |
| ➤ 600 ++ Concept Wise Tests | ➤ A Strong Subject Foundation |
| ➤ 50 ++ Chapter Wise Tests | ➤ A Refined Learning Methodology |
| ➤ 50 ++ Model Tests | ➤ Updated Study materials |
| ➤ 2000 ++ Problem Discussions | ➤ Freshers Can easily understand |
| ➤ Recorded Videos | ➤ Question banks |




Q.21 – Q.60 MCQ, carry TWO marks each (for each wrong answer: – 0.5).

You are required to Answer Maximum 35 Questions.

21. Dinuclear anion $[\text{I}_2(\text{OH})_2\text{O}_8]^{4-}$ has two bridging oxo groups. The geometry around each iodine is
 (a) octahedral (b) monocapped octahedral
 (c) square pyramidal (d) pentagonal bipyramidal
22. Using a double beam UV-visible spectrophotometer, Beer's law fails for $\text{K}_2\text{Cr}_2\text{O}_7$ solution when
 (a) intensity of light source is changed (b) detector is not a photomultiplier tube
 (c) cuvette of 2 cm size is used (d) pH is not kept same in all measurements
23. Trivalent lanthanide ion having isotropic magnetic susceptibility is
 (a) Eu^{3+} (b) Gd^{3+} (c) Yb^{3+} (d) Lu^{3+}
24. The structure of CaB_6 is close to that of
 (a) cesium chloride (b) nickel arsenide (c) rock salt (d) zinc blende
25. The correct order of C–O bond length is
 (a) $\text{H}_3\text{B.CO} > [\text{Mn}(\text{CO})_6]^+ > [\text{Cr}(\text{CO})_6]^- > [\text{V}(\text{CO})_6]^-$
 (b) $[\text{V}(\text{CO})_6]^- > [\text{Cr}(\text{CO})_6]^- > [\text{Mn}(\text{CO})_6]^+ > \text{H}_3\text{B.CO}$
 (c) $[\text{Mn}(\text{CO})_6]^+ > \text{H}_3\text{B.CO} > [\text{V}(\text{CO})_6]^- > [\text{Cr}(\text{CO})_6]^-$
 (d) $[\text{Cr}(\text{CO})_6]^- > [\text{V}(\text{CO})_6]^- > \text{H}_3\text{B.CO} > [\text{Mn}(\text{CO})_6]^+$
26. Among the elements Zn, Ga, Ge and As, the one with the lowest first ionization energy is
 (a) As (b) Zn (c) Ga (d) Ge
27. The total degeneracy of the ground term of Co^{II} (high spin) in octahedral geometry is
 (a) 18 (b) 12 (c) 28 (d) 9
28. For the following reaction, the structure of the intermediate is:





29. High spin complex of a 3d metal ion M has a magnetic moment of 2.9 B.M in octahedral coordination environment and 4.1 B.M in tetrahedral environment. The M ion is

- (a) Co^{III} (b) Ni^{II} (c) Cu^{II} (d) Co^{II}

30. For electronic spectra of $\text{K}_2\text{Cr}_2\text{O}_4$ (X) and K_2MoO_4 (Y) the correct combination is

- (a) transition is d-d and λ_{max} for X < Y (b) transition is LMCT and λ_{max} for X < Y
(c) transition is LMCT and λ_{max} for X > Y (d) transition is MLCT and λ_{max} for X > Y

31. Removal of an electron from NO molecule results in

- [P] an increase in the $\nu_{(\text{NO})}$ in the IR spectrum
[Q] an EPR active species
[R] electrons in HOMOs being closer to the oxygen than to nitrogen 2p orbitals
[S] electrons in HOMOs being closer to the nitrogen than to oxygen 2p orbitals

The correct answer is

- (a) P only (b) P and R (c) Q and S (d) P, Q, and R

32. Consider the nature of solvents in column-I and the corresponding λ_{max} for I_2 in various solvents given in column-II. (for I_2 vapor λ_{max} is 520 nm). Match Column-I with Column-II

	Column I	Column II (λ_{max} , nm)	P	Q	R	S
(P)	Non-donor (i)	520	(a) i ; ii ; iii ; iv			
(Q)	Weak donor (ii)	500	(b) iii ; iv ; ii ; i			
(R)	Strong donor (iii)	450	(c) i ; iii ; iv ; ii			
(S)	π electron donor (iv)	360	(d) iv ; iii ; ii ; i			

33. For the catalytic activity of Cu and Zn containing enzyme, superoxide dismutase, what is/are the correct statement(s)?

- [P] Cu and Zn both are essential
[Q] only Cu is essential
[R] Zn is essential and Cu may be replaced by any other divalent metal atom

[S] Zn may be replaced by any other divalent metal atom


- (a) P only (b) R only (c) S only (d) Q and S

34. Mass spectrum of a compound shows an $[M + 2]$ ion peak that is about 4% of M^+ .

This indicates that the compound has one

- (a) fluorine (b) sulfur (c) bromine (d) Chlorine


35. The major product formed in the following reaction is




36. For the following compounds, the correct order of reactivity towards nucleophilic acyl substitution is

- (a) acetyl chloride < methyl acetate < Ac_2O < acetamide
 (b) acetamide < methyl acetate < Ac_2O < acetyl chloride
 (c) acetamide < Ac_2O < acetyl chloride < methyl acetate
 (d) methyl acetate < acetamide < Ac_2O < acetyl chloride


37. The major product formed in the following reaction is:




38. The major product formed in the following reaction is:




39. The **major product** formed in the following reaction is



40. The **major product** formed in the following reaction is



41. **Number of signals** observed in the $^{13}\text{C-NMR}$ spectrum of the following compound is





(a) 4



(b) 5

(c) 6




(d) 8

42. The structure of the product formed during the reaction of **amino acid** with **ninhydrin** is





43. The correct order of rate of solvolysis in 80% ethanol at 25 °C is



(a) Q > R > P

(b) P > Q > R

(c) R > Q > P

(d) R > P > Q

44. IUPAC nomenclature of following propellane is





(a) tricyclo[1.1.1.0^{2,4}]pentane

(b) tricyclo[1.1.0.1^{1,3}]pentane

(c) tricyclo[1.1.1^{1,3}.0^{1,5}]pentane

(d) tricyclo[1.1.1.0^{1,3}]pentane

45. The correct statement about following species is





(a) Both I and II are aromatic

(b) I is aromatic and II is antiaromatic

(c) I is nonaromatic and II is antiaromatic

(d) I is aromatic and II is homoaromatic

46. The correct statement about the following compounds is



(a) I is more stable than II

(b) II is more stable than I

(c) I and II are equally stable

(d) I and II are both locked conformations

47. In the pure Raman rotational spectrum of $^{16}\text{O}_2$, whose electronic ground state is $^3\Sigma_g^-$, transitions between to/from


(a) even J levels are missing

(b) odd J levels are missing

(c) all J levels appear

(d) none of the J levels appear

48. Elementary steps of a reaction are as follows



If steady state approximation is applicable to C, the rate of product formation in the reaction is

(a) proportional to $[A][B]$ (b) proportional to $[A]^2[B]^2$ (c) proportional to $[A]^{1/2}[B]^{1/2}$ (d) independent of $[A][B]$ 49. The term symbol for the ground state of B_2 , is(a) $^1\Sigma_g^+$ (b) Σ_g^- (c) $^3\Sigma_g^-$ (d) $^3\Sigma_g^+$

50. A Gaussian distribution has the functional form $f(x) = \frac{2}{\sqrt{2a^2\pi}} e^{-(x-b)^2/2a^2}$. The variance of such distribution is

(a) a

(b) a^2

(c) b

(d) b^2

51. The change in entropy for a reversible adiabatic process is

(a) maximum

(b) Minimum

(c) zero

(d) Positive

52. The standard cell potential for the reaction

$Zn_{(s)} + Cu^{2+}_{(aq)} \rightleftharpoons Zn^{2+}_{(aq)} + Cu_{(s)}$ is +1.10 V. The Gibbs free energy change during the reaction is (F = 96500 coulomb mol⁻¹)

(a) -21.2 kJ mol⁻¹(b) +212 kJ mol⁻¹(c) -212 kJ mol⁻¹(d) -212 J mol⁻¹

53. If the unit of the rate constant of a reaction is L³ mol⁻³ s⁻¹, the order of the reaction is

(a) 1

(b) 2

(c) 3

(d) 4

54. The lowest energy state of a $1s^1 2s^1$ electronic configuration, according to Hund's rule, is,

(a) 3S_0 (b) 1S_0 (c) 3S_1 (d) 1S_1

55. The commutator of $\hat{x}$ with the Hamiltonian $\hat{H}$, $[\hat{x}, \hat{H}]$, is


(a) 0

(b) $i\hbar$ (c) $\frac{-\hbar^2}{2m} \hat{p}_x$ (d) $\frac{i\hbar}{m} \hat{p}_x$

56. A 5 g/L polymer solution is prepared with a polymer whose molar mass is 25 kg.

The osmotic pressure (in atm) of this solution at 25 °C is ____ (RT ≈ 2500 J mol⁻¹)

- (a) 0.002 (b) 0.05 (c) 0.005 (d) 0.008
57. If all the lattice points of an **FCC** structure are occupied by uniform hard spheres that touch each other, the **fraction of volume occupied** is
- (a) $\frac{\pi\sqrt{2}}{6}$ (b) $\frac{\pi\sqrt{3}}{6}$ (c) $\frac{\pi}{6}$ (d) $\frac{2\pi}{6}$
58. **Origin of the colligative properties** of a dilute solution is
- (a) volatility of solute molecule (b) interaction of solute-solvent molecules
(c) zero enthalpy of mixing (d) entropy of mixing
59. The graph that represents the **Langmuir Adsorption Isotherm** is




60. Correct match for the **coenzymes in Column-I** with their **function in Column-II** is


	Column-I	Column-II	P	Q	R
P.	NaDH	i. Oxidation	(a)	i ; ii ; iii	
Q.	FAD	ii. Acyl group transfer	(b)	iii ; i ; ii	
R.	CoASH	iii. Reduction	(c)	iii ; ii ; i	
			(d)	ii ; i ; iii	

Q.61 – Q.120 MCQ, carry FOUR marks each (for each wrong answer: – 1).

You are required to Answer Maximum 25 Questions.

61. One of the products formed in the **bis(η^3 -allyl)nickel** complex catalyzed cyclo-dimerization of butadiene in the presence of PR_3 is compound-X given below. **Identify its precursor.**





62. The transformation are given in column-I and reagent in column-II. Match the items of column-I with those of column-II.

	Column-I		Column-II
[P]	$[\text{MnO}_4]^- \rightarrow [\text{MnO}_4]^{2-}$	I.	H_2SO_4
[Q]	$\text{Me}_3\text{CH} \rightarrow [\text{Me}_3\text{C}]^+$	II.	Na in liquid NH_3
[R]	$\text{Ag} + \text{Au} \rightarrow \text{Ag}[\text{AuF}_4]$	III.	$[\text{H}_2\text{SO}_3\text{F}]^+$ (super acid)
[S]	$\text{H}_3\text{PO}_4 \rightarrow [\text{P}(\text{OH})_4]^+$	IV.	Liquid BrF_3

The correct match is

	P	Q	R	S
(a)	I ; II	III ; IV		
(b)	II ; III	IV ; I		
(c)	III ; II	I ; IV		
(d)	III ; I	IV ; II		

63. Consider the following statements for the oxygenation of hemocyanine:

[P] oxidation state of both copper atoms changes by two

[Q] it becomes intense blue from colourless

[R] dioxygen is reduced to O_2^{2-}

[S] the $\mu\text{-}\eta^2\text{:}\eta^2$ bond forms between each oxygen and copper atoms


The correct statements are:

- (a) P and R (b) Q and R (c) P, Q and R (d) Q, R and S

64. The correct increasing order of C-C bond length in the following molecules (P-R) is

P	Q	R	
$[\text{PtCl}_3(\text{C}_2\text{H}_4)]^-$	$[\text{Pt}(\text{PPh}_3)_2(\text{C}_2\text{H}_4)]$	$[\text{Pt}(\text{PPh}_3)_2\{\text{C}_2(\text{CN})_4\}]$	
(a) $\text{R} < \text{P} < \text{Q}$	(b) $\text{P} < \text{Q} < \text{R}$	(c) $\text{Q} < \text{R} < \text{P}$	(d) $\text{R} < \text{Q} < \text{P}$

65. Which of the following are NOT closo clusters?





The correct answer is


- (a) R and S (b) P and Q (c) P and R (d) Q and R

66. Identify the pair of molecules which are isoelectronic as well as isostructural from the following:



- (a) Q and R (b) P and U (c) P and S (d) R and T

67. For the following reaction, correct statement(s) is/are



[P] Oxidation State of iridium increases from I to III

[Q] It is β -hydride elimination reaction


[R] (I) and (II) both are diamagnetic

[S] It is migratory insertion reaction

The correct answer is

- (a) P only (b) P and R (c) R and S (d) Q, R and S

68. The reaction of $[(\eta^5\text{-C}_5\text{H}_5)\text{Fe}(\text{CO})_2]^-$ with CH_3I gives compound-X. The $^1\text{H-NMR}$ spectrum of X shows two singlets in an integrated intensity ratio of 3 : 5. Compound-X upon reaction with PPh_3 gives compound-Y. The $^1\text{H-NMR}$ spectrum of Y shows 3 sets of signals in an integrated intensity ratio of 3 : 5 : 15. Compounds X and Y respectively, are:



69. Identify the correct statements about the electronegativity of groups given below:

- [P] CF_3 group has greater value than that of NF_2
 [Q] NH_2 group has lower value than that of NF_2
 [R] OH group has greater value than that of NF_2
 [S] CH_3 and C_2H_5 groups have almost similar values

Correct answer is

- (a) P, Q and S (b) Q and R (c) Q, R and S (d) Q and S

70. Height equivalent to theoretical plate (HETP) in gas-liquid chromatography depends significantly on which of the following the Correct answer is

P

Temperature of column

Q

Velocity of carrier gas

R

Packing of column

S

Column material

- (a) P, Q and R (b) R and S (c) Q, R and S (d) P and R

71. A **binary fluoride (Z)** of xenon combines with two moles of NaF to give a product which on heating to 100 °C affords compound-P. The **alkaline hydrolysis** of P gives **perxenate salt**. **Z** and **P** are, respectively,

(a) XeF_2 and XeF_4 (b) XeF_4 and XeF_6 (c) XeF_6 and XeF_4 (d) XeF_6 and XeF_6

72. Match **fluorescence colour** given in column-I with lanthanide ions given in column-II

	Column-I	Column-II	i	ii	iii	iv
(i)	Pink	(P)	Sm(III)	(a) P ; R ; Q ; S		
(ii)	Red	(Q)	Tb(III)	(b) S ; R ; Q ; P		
(iii)	Green	(R)	Eu(III)	(c) P ; Q ; R ; S		
(iv)	Blue	(S)	Tm(III)	(d) R ; Q ; S ; P		

73. Choose the correct set of statements for **cisplatin**,

[P] It can be prepared from $\text{K}_2[\text{PtCl}_4]$

[Q] It can be prepared from $[\text{Pt}(\text{NH}_3)_4]\text{Cl}_2$

[R] In its preparation, the observed trans effect for Cl^- is greater than that of NH_3

[S] In blood it stays in equilibrium with $\text{cis-}[\text{Pt}(\text{NH}_3)_2\text{Cl}(\text{H}_2\text{O})]^+$

[T] In DNA strand, it binds to two adjacent cytosine bases

The correct set is

(a) P, R and S (b) P, R, S and T (c) Q, R and S (d) Q, R, S and T

74. In fission of ^{235}U atom the energy released is **200 MeV**. In one day fission of **1 kg ^{235}U** will give power (in MW) approximately

(a) 550 (b) 650 (c) 950 (d) 1250

75. In the structures



P and Q are made up of two MCl_4 units. For these structures, which statement is correct?

- (a) P and Q both have MCl_4 units eclipsed
 (b) P and Q both have MCl_4 units staggered
 (c) P has both MCl_4 units staggered and Q has both MCl_4 units eclipsed
 (d) P has both MCl_4 units eclipsed and Q has both MCl_4 units staggered

76. For the Wacker process, pick the correct statement(s) from the following:

[P] Pd(II) is reduced to Pd(0) by Cu(I)

[Q] Pd(0) is oxidized to Pd(II) by Cu(II)

[R] Cu(II) promotes the reductive elimination

Correct answer is

- (a) P and R (b) Q and R (c) P and Q (d) Q only

77. Consider following statements


I. AsCl₅ is thermally less stable than PCl₅

II. Size of As is more than that of P

Choose correct answer from the following

- (a) Statement I and II are true and II is the correct explanation of I
 (b) Statements I and II are true but II is not the correct explanation for I
 (c) Statement I is true and statement II is false
 (d) Both the statements I and II are false

78. Consider the following statements for



[P] There is an M-M (M = Be, B, Ga) bond in all

[Q] The oxidation state of Be, B and Ga is +2

[R] The geometry around the central atom is planar for all

[S] The geometry around the central atom is planar in I and II only

The correct statement(s) is/are

- (a) P, Q and R (b) P and Q (c) S only (d) Q, R and S

79. Consider the following statements:

[P] Cr²⁺ is easier to oxidise than V²⁺ in the gas phase


[Q] Cr²⁺_(aq) is a more powerful reducing agent than V²⁺_(aq)

[R] The rate of water exchange for Cr²⁺_(aq) (aq) is much faster than for V²⁺_(aq)

The correct statements are

- (a) P and Q (b) P and R (c) Q and R (d) P, Q and R

80. Consider the statements P-S regarding equations I- III:



[P] Marcus equation is applicable to I and II

[Q] Marcus equation is applicable to II only


[R] Equations I and II involve inner sphere electron transfer

[S] Equations I and III involve inner sphere electron transfer


The correct statements are:

- (a) P and Q (b) Q and R (c) Q and S (d) R and S


81. The major product formed in the following oxidation reaction is:




82. The major product formed in the following reaction is




83. The major products P and Q formed in the following reaction sequence are








84. The major products **P** and **Q** formed in the following reaction sequence are



85. Structure of the **intermediate-X** and the final **product-Y** in the following reaction sequence are, (dba = dibenzylidene acetone)



86. Mechanism of the following transformation involves



- (a) A [2+2] cycloaddition followed by 'con' rotatory electrocyclic ring opening
 (b) A [4+2] cycloaddition followed by 'con' rotatory electrocyclic ring opening
 (c) A [4+2] cycloaddition followed by Cope rearrangement
 (d) A [2+2] cycloaddition followed by 'dis' rotatory electrocyclic ring opening

87. The correct statement about solvolysis using NaOAc/AcOH of following compounds is




- (a) P reacts faster than Q to give trans-1,2-diacetoxycyclohexane
 (b) Q reacts faster than P to give trans-1,2-diacetoxycyclohexane

(c) P reacts faster than Q to give cis-1,2-diacetoxycyclohexane

(d) Q reacts faster than P to give cis-1,2-diacetoxycyclohexane

88. The structure of the intermediate-X and the major product-Y, formed in the following reaction are



89. A compound shows following spectral data:


¹H-NMR : 7.9 (d, J = 8 Hz, 2H), 6.6 (d, J = 8 Hz, 2H), 4.3 (q, J = 6 Hz, 2H), 4.0 (br, s, 2H, D₂O exchangeable), 1.4 (t, J = 6 Hz, 3H)


Mass : m/z 165, 137, 120, 92

The correct structure of the compound is




90. The major allylic alcohol-P and the ester-Q formed in the following reaction sequence are






91. The major products **P** and **Q** formed in the following reaction sequence are




92. Reaction of **D-glucose** with following reagents produces

Reagents : i. Acetone/ H^+
 ii. PDC
 iii. $(EtO)_2P(O)CH_2CO_2Et$, NaH




93. For the following thermal [2+2] cycloaddition reaction, the correct statement about Transition State(TS) and preference for endo product formation is




- (a) $\pi_{2s} + \pi_{2s}$; Me > i-Pr > t-Bu
 (b) $\pi_{2s} + \pi_{2a}$; t-Bu > i-Pr > Me
 (c) $\pi_{2s} + \pi_{2a}$; Me > i-Pr > t-Bu
 (d) $\pi_{2s} + \pi_{2s}$; t-Bu > i-Pr > Me

94. The major products **P** and **Q** formed in the following reaction sequence are:



- (a) ; (b) ;
 (c) ; (d) ;

95. The major products formed in the following photochemical reaction are




- (a) + (b) +
 (c) + (d) +


96. Irradiation of either **cis-** or **trans-stilbene** at 313 nm results in the formation of a mixture of **93 % cis** and **7 % trans** olefin because

- (a) trans-stilbene is more stable than cis-stilbene
 (b) the extinction coefficient of trans-stilbene is greater than cis-stilbene at exciting wavelength
 (c) the transition state structures of cis and trans stilbenes are different
 (d) the triplet excited states of cis and trans stilbenes are at different energy levels


97. The major heterocyclic compound formed in the following reaction is



98. The major products **P** and **Q** formed in the following reaction sequence are



99. The correct order of rates for the following reactions is




(a) $k_1 > k_2$ and $k_3 > k_4$

(b) $k_1 > k_2$ and $k_4 > k_3$

(c) $k_2 > k_1$ and $k_3 > k_4$

(d) $k_2 > k_1$ and $k_4 > k_3$

100. The correct match of protons in Column-I with the $^1\text{H-NMR}$ chemical shifts in Column-II for the product of the following reaction is



	Column-I	Column-II (δ ppm)
P.	H_A	i. -0.3
Q.	H_B	ii. 5.1
R.	$H_{1 \& 7}$	iii. 6.4
S.	H_{2-6}	iv. 8.5

	P	Q	R	S
(a)	ii	i	iii	iv
(b)	i	ii	iv	iii
(c)	iv	i	iii	ii
(d)	ii	iv	i	iii

101. Which of these is **not a suitable unnormalized wave function** for the excited $1s^1 2s^1$ electron configuration of the helium atom?

- (a) $[1s(1)2s(2) - 2s(1)1s(2)][\beta(1)\beta(2)]$
 (b) $[1s(1)2s(2) + 2s(1)1s(2)][\alpha(1)\beta(2) - \beta(1)\alpha(2)]$
 (c) $[1s(1)2s(2) - 2s(1)1s(2)][\alpha(1)\beta(2) + \beta(1)\alpha(2)]$
 (d) $[1s(1)2s(2) + 2s(1)1s(2)][\alpha(1)\beta(2)]$

102. Two opposite sides (**in the y-direction**) of a square box of side L are slightly stretched. Consider the following four statements:

- [P] The point group changes from D_{4h} to D_{2h}
 [Q] The (1,2) and (2,1) energy levels remain doubly degenerate
 [R] Both the energy levels are lowered and the energy of the (1,2) level is higher than that of the (2,1) level
 [S] Both the energy levels are lowered and the energy of the (1,2) level is lower than that of the (2,1) level

The two correct statements are:

- (a) P and Q (b) P and R (c) Q and R (d) P and S
103. Consider a model system of five non-interacting fermions in a single 3-dimensional harmonic oscillator. The Hamiltonian of a single particle is

$$\hat{H} = \frac{1}{2m} (\hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2) + \frac{1}{2} m \omega^2 (x^2 + y^2 + z^2)$$

Where m is the mass of the particle, ω is the angular frequency, $\hat{p}_x$, $\hat{p}_y$ and $\hat{p}_z$ are the momentum operators. The ground state energy of the system of 5 non-interacting fermions is

- (a) $\frac{21}{2} \hbar \omega$ (b) $\frac{15}{2} \hbar \omega$ (c) $\frac{5}{2} \hbar \omega$ (d) $\frac{25}{2} \hbar \omega$

104. A particle is in a state $\phi = \varphi_1 + 3\varphi_2$, where φ_1 and φ_2 are eigenfunctions of the Hamiltonian of the particle with eigenvalues E_1 and E_2 , respectively. The average energy of the particle in the state ϕ is

- (a) $(E_1 + 9E_2)/10$ (b) $(E_1 + 3E_2)$ (c) $(E_1 + 9E_2)/4$ (d) $(E_1 + 3E_2)/10$


105. Which of the following statements on ground state perturbation theory, involving the zeroth order energy $E_0^{(0)}$, first order energy correction $E_0^{(1)}$ and second order energy correction $E_0^{(2)}$, is false?

- (a) $E_0^{(1)}$ is the average value of perturbation operator with respect to the ground state of the zeroth order Hamiltonian
 (b) $E_0^{(1)}$ is necessarily negative
 (c) $E_0^{(2)}$ is necessarily negative
 (d) $E_0^{(0)} + E_0^{(1)}$ is an upper bound to the exact ground state energy

106. Difference between activation energies of the reverse and forward steps of a reversible reaction is $9.212 RT$. If the pre-exponential factor of the forward reaction is double that of the reverse reaction at the same temperature, the equilibrium constant for the reaction at that temperature will be $(\ln 10 = 2.303)$

- (a) 1×10^4 (b) 2×10^4 (c) 1×10^{-4} (d) 2×10^{-4}

107. For an enzyme-substrate reaction,



the slope and the intercept of the plot between $\frac{1}{r}$ and $\frac{1}{[S]}$ are $10^{-2} s$ and $10^2 M^{-1} s$, respectively. If $E_0 = 10^{-6} M$ and $\frac{k_{-1}}{k_2} = 1000$, the value of k_1 (in units of $M^{-1}s^{-1}$) will be close to,

[r is the rate of the reaction and E_0 is the initial concentration of the enzyme]

- (a) 1×10^{11} (b) 1×10^4 (c) 1×10^8 (d) 1×10^6



108. Translational partition function of a D_2 molecule confined in a 100 cm^3 vessel at 25°C is, $(h = 6.626 \times 10^{-34}\text{ J s}, k = 1.381 \times 10^{-23}\text{ J K}^{-1})$

(a) 3.8×10^{22} (b) 5.8×10^{24} (c) 7.8×10^{26} (d) 9.8×10^{28}

109. The volume (cm^3) of CO adsorbed on charcoal (273 K) at two different pressures is given below

P(kPa)	40	80
V(cm^3)	25	40

Assuming Langmuir isotherm, the maximum possible volume (cm^3) CO that can be adsorbed is

(a) 50 (b) 100 (c) 150 (d) 200

110. The number of lines in EPR spectrum of CD_3 ($I_D = 1$) is

(a) 3 (b) 5 (c) 7 (d) 9

111. A symmetric top molecule, among the following is

(a) ethylene (b) Allene (c) butatriene (d) Hexatriene

112. The allowed electronic transition in fluorine molecule is

(a) $\Sigma_g^+ \rightarrow \Sigma_g^+$ (b) $\Sigma_g^+ \rightarrow \Sigma_g^-$ (c) $\Sigma_g^+ \rightarrow \Pi_u$ (d) $\Sigma_g^+ \rightarrow \Delta_u$

113. Assuming harmonic approximation, the energy change for the reaction $HCl + D_2 \rightarrow DCl + HD$ in cm^{-1} is,

(the vibrational frequency data in cm^{-1} is given in the table below),

HCl	D_2	DCl	HD
2885	2990	1990	3627

(a) -258 (b) +258 (c) -129 (d) +129

114. The transition moment integral for a rotational transition between $J = 1; M_J = 0$ and $J = 2; M_J = 0$ states for a diatomic molecule along the z axis is proportional to

(a) $\int_0^\pi \cos^2\theta(3\cos^2\theta - 1)d\theta$ (b) $\int_0^\pi \cos^2\theta(3\cos^2\theta - 1)\sin\theta d\theta$

(c) $\int_0^\pi \cos\theta(3\cos^2\theta - 1)\sin\theta d\theta$ (d) $\int_0^\pi \cos\theta(3\cos^2\theta - 1)\sin^2\theta d\theta$

115. One of the correct normalized sp^2 hybrid orbitals is

(a) $\frac{1}{3}\phi_{2s} + \frac{1}{3}\phi_{2p_x} + \frac{1}{3}\phi_{2p_y}$ (b) $\frac{1}{2}\phi_{2s} + \frac{\sqrt{3}}{\sqrt{8}}\phi_{2p_x} + \frac{\sqrt{3}}{\sqrt{8}}\phi_{2p_y}$

(c) $\frac{1}{\sqrt{3}}\phi_{2s} + \frac{\sqrt{2}}{\sqrt{3}}\phi_{2p_x}$ (d) $\frac{1}{3}\phi_{2s} + \frac{2}{3}\phi_{2p_x}$

116. At 300 K, the **thermal expansion coefficient** and the isothermal compressibility of liquid water are $2 \times 10^{-4} \text{ K}^{-1}$ and $5 \times 10^{-5} \text{ bar}^{-1}$, respectively. $\left(\frac{\partial U}{\partial V}\right)_T$ (in k bar) for water at 320 K and 1 bar will be
 (a) 2.4 (b) 1.2 (c) 0.6 (d) 12.0
117. In the **phase diagram of water**, the **solid liquid boundary has a negative slope**. The reason for this unusual behaviour can be traced to decrease in
 (a) density of the system on melting (b) volume of the system on melting
 (c) entropy of the system on melting (d) enthalpy of the system on melting
118. The **standard cell potential of cell**, $\text{Pt} | \text{H}_{2(g)} | \text{HBr}_{(aq)} | \text{AgBr}_{(s)} | \text{Ag}_{(s)}$, was measured over a range of temperatures, and the data was fitted as

$$E^0(\text{Volt}) = 0.01 - 1 \times 10^{-4}(T - 298) - 2 \times 10^{-6}(T - 298)^2$$
 The **standard reaction entropy ($\text{JK}^{-1}\text{mol}^{-1}$)** and **enthalpy (kJmol^{-1})** at 298 K are
 (a) -9.65 and -3.84 (b) -3.84 and -9.65 (c) -18.3 and -7.68 (d) -7.68 and -18.3
119. The (002) plane of an **elemental FCC crystal** diffracts X-rays ($\lambda = 0.154 \text{ nm}$) at **Bragg angle 90°** . The density of the crystal is $4 \times 10^4 \text{ kg m}^{-3}$. The **atomic weight of the elemental solid is**
 (a) 22 (b) 44 (c) 88 (d) 66
120. A solution of Fe^{3+} is titrated **potentiometrically** using Ce^{3+} solution at 25°C . The emf (in V) of the redox system thus formed when,
 (i) 50 % of Fe^{3+} and (ii) 80 % of Fe^{3+} are titrated, would respectively be
 (Given $E^0_{\text{Fe}^{3+}/\text{Fe}^{2+}} = 0.77\text{V}$, $\log_{10} 2 = 0.301$)
 (a) 0.734 and 0.77 (b) 0.77 and 0.385 (c) 0.77 and 0.734 (d) 0.385 and 0.367

Answer Key

PART - B

Q.No	Ans
21.	a
22.	d
23.	b

Q.No	Ans
31.	b
32.	c
33.	d

Q.No	Ans
41.	c
42.	a
43.	d

Q.No	Ans
51.	c
52.	c
53.	d

24.	a
25.	b
26.	c
27.	c
28.	d
29.	b
30.	c

34.	b
35.	a
36.	b
37.	d
38.	b
39.	c
40.	a

44.	d
45.	b
46.	c
47.	a
48.	a
49.	c
50.	b

54.	c
55.	d
56.	c
57.	a
58.	d
59.	d
60.	b

PART - C

Q.No	Ans
61.	b
62.	b
63.	d
64.	b
65.	d
66.	b
67.	b
68.	b
69.	d
70.	a
71.	d
72.	a
73.	a
74.	c
75.	d

Q.No	Ans
76.	d
77.	b
78.	c
79.	c
80.	c
81.	b
82.	a
83.	a
84.	c
85.	a
86.	d
87.	a
88.	a
89.	a
90.	b

Q.No	Ans
91.	d
92.	a
93.	b
94.	a
95.	a
96.	b
97.	c
98.	a
99.	b
100.	a
101.	d
102.	d
103.	a
104.	a
105.	b

Q.No	Ans
106.	b
107.	a
108.	c
109.	b
110.	c
111.	b
112.	c
113.	c
114.	b
115.	c
116.	b
117.	b
118.	a
119.	a
120.	c



- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © Inspite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So, we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.

- © இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

