

# GATE – 2001 – Chemistry



#### www.csircoaching.com

- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

Featu	res
> 300 ++ Live Classes	A Well-Defined Curriculum
200 ++ Concept Wise Tests	<ul> <li>A Strong Subject Foundation</li> </ul>
$\succ$ 50 ++ Chapter Wise Tests	> A Refined Learning Methodology
$\succ$ 50 ++ Model Tests	<ul><li>Updated Study materials</li></ul>
> 2000 ++ Problem Discussions	<ul> <li>Freshers Can easily understand</li> </ul>
<ul><li>Recorded Videos</li></ul>	$\succ$ Question banks

 $\bigoplus$ 



	Q.1 – Q.50 Multiple Choice Question (MCQ), carry TWO marks each								
	<u>(for each wrong answer: – 2/3).</u>								
1.	Icosahedral structure is generally exhibited by								
	(a) C	(b) Si	(c) Ge	(d) B					
2.	The hybrid orbital	s used by bromine at	om in BrF <sub>3</sub> are						
	(a) sp <sup>2</sup>	(b) sp <sup>3</sup>	(c) $sp^3d$	(d) $sp^3d^2$					
3.	The metal ion pres	ent in carbonic anhy	drase is						
	(a) Mn	(b) Zn	(c) Cu	(d) Fe					
4.	The most acidic ac	ua ion is							
	(a) $[Fe(H_20)_6]^{3+}$	(b) [Co(H <sub>2</sub> O) <sub>6</sub> ] <sup>3+</sup>	(c) $[Ti(H_20)_6]^{3+}$	(d) $[Cr(H_2O)_6]^{3+}$					
5.	Which one of the f	ollowing metal fragm	ents, d <sup>n</sup> - ML <sub>m</sub> , is iso	lobal with CH?					
	(a) $d^{7}$ -ML <sub>5</sub>	(b) $d^8$ -ML <sub>4</sub>	(c) $d^9-ML_3$	(d) $d^{5}$ -ML <sub>6</sub>					
6.	The softest acid an	nongst the following i	s						
	(a) Al <sup>3+</sup>	(b) Li <sup>+</sup>	(c) $Ca^{2+}$	(d) Ag <sup>+</sup>					
7.	The chromium	(III) species forme	d soon after elec	tron transfer between					
	[IrCl <sub>6</sub> ] <sup>2-</sup> and [Cr(	$(H_2 0)_6]^{2+}$ is :							
	(a) $[Cr(H_20)_6]^{3+}$	(b) $[Cr(H_2O)_5Cl]^2$	+ (c) $[CrCl_6]^{3-}$	(d) $[Cr(H_2O)_3Cl_3]$					
8.	The strongest redu	icing ion of the follow	ving is						
	(a) U <sup>3+</sup>	(b) Am <sup>3+</sup>	(c) Cm <sup>3+</sup>	(d) Cf <sup>3+</sup>					
9.	The first ionization	n potential of Mg, Al,	P and S follows the o	order					
	(a) $Mg < Al < P <$	s <b>CM</b>	(b) Al < Mg <	P < S					
	(c) $Al < Mg < S <$	Р	(d) Mg < Al <	S < P					
10.	As per the uncerta	inty principle, Δx. Δp	$y_y =$						
	(a) h	(b) h/2π	(c) λ	(d) zero					
11.	The second lower s	state of <mark>particle in a</mark> c	ubic box is						
	(a) non degenerate	(b) doubly degenerate	(c) triply degenerate	(d) six-fold degenerate					
12.	In comparison to t	he frequency of EPR	transition and NMR	transition frequency is					
	(a) much higher	(b) much lower	(c) almost same	(d) none of these					
13.	The symmetry poi	nt group of the BF <sub>3</sub> n	nolecule is:						
	(a) C <sub>3v</sub>	(b) D <sub>3h</sub>	(c) C <sub>2v</sub>	(d) D <sub>2h</sub>					
14.	For an irreversibl	e adiabatic expansio	n of a perfect gas f	rom volume $V_i$ to $V_f$ the					
	change in entropy of the gas is:								

 $\bigoplus$ 





1

2

	(a) nR ln ( $V_f/V_i$ ) (b) Zero (c) less than zero (d) greater the formula of the second seco	nan zero
15.	5. For the cell Ag (s)   AgCl (satd), NaCl $(aq,m_1)$   NaCl $(aq,m_2)$ , AgCl (sat	d)   Ag(s)
	where $m_1$ and $m_2$ are different molalities ( $m_1 > m_2$ ), the standard cell pot	ential is:
	(a) $-RT \ln (m_1/m_2)$ (b) Zero (c) $-RT \ln (a_1/a_2)$ (d) $-F_1$	RT ln (a <sub>1</sub> )
16.	6. For an ideal dilute solution, which one of the following statements is correc	t?
	( $\gamma$ and $\chi$ are activity coefficient and mole fraction res	pectively)
	(a) $\gamma$ (solvent) $\rightarrow 1$ as $\chi$ (solvent) $\rightarrow 0$ (b) $\gamma$ (solute) $\rightarrow 1$ as $\chi$ (solute)	$\rightarrow 1$
	(c) $\gamma$ (solvent) $\rightarrow 1$ as $\chi$ (solvent) $\rightarrow 1$ (d) $\gamma$ (solvent) $\rightarrow 0$ as $\chi$ (solvent)	$(t) \rightarrow 1$
17.	7. For the reaction: $Br_2(g) + BF_2(g) \rightarrow 2BrF_3(g)$ , the equilibrium constant	at 2000 K
	and 1.0 bar is 5.25. When the pressure is increased by 8-fold, the eq	uilibrium
	constant,	
	(a) Increase by a factor of 1.86 (b) Decreases by a factor of 1.86	5
	(c) Remains same (d) Increases by a factor of 8	
18.	8. 2-phenylethanol may be prepared by the reaction of phenyl magnesium brow	mide with
	(a) HCHO (b) $CH_3CHO$ (c) $CH_3COCH_3$ (d) $\overset{O}{\rightharpoonup}$	
19.	9. o-chlorotoluene reacts with sodamide in liquid ammonia to give o-toluidin	e, and m-
	toluidine. This reaction proceeds through an intermediate	
	(a) $CH_3$ (b) $CH_3$ (c) $CH_3$ (c) $CH_3$ (d) $CH_3$	l <sub>3</sub>
		CI
20.	20. The number of signals observed in <sup>1</sup> H-NMR spectrum of 3,5-dibromotoluer	ne is:
	(a) 3 (b) 4 (c) 2 (d) 6	
21.	21. Which one of the following molecules will have $n \to \pi^*$ transition at the	e longest
	wavelength?	
	(a) HCHO (b) $CH_3COC_2H_5$ (c) $C_6H_5COC_6H_5$ (d) $CH_5$	<sub>3</sub> COC <sub>6</sub> H <sub>5</sub>
22.	2. The reaction of cyclooctyne with $HgSO_4$ in the presence of aqueous $H_2SO_4$	gives
	(a) OH (b) OH (c) (d)	<u>_0</u>
	С ОН С	
23.	$\begin{array}{ccc} & & & \Theta \\ & & & \Theta \\ & & & & \\ & & & & \\ & & & &$	
	$\frac{\text{Pyrolysis of }}{\text{Me}} \frac{\text{Me}}{\text{CD}_3} = \frac{\text{Would be}}{\text{Me}}$	
	(a) A mixture of $H_2C=CH-CD_3$ and $H_3C-CH=CD_2$ (b) $H_3C-CH=CD_3$	2
Tiruc	iruchirannalli – 620.024 www.csircoaching.com	Damail com
muc	www.csi.couching.com	

**Q** 

(c) 
$$\bigotimes_{M=2}^{\oplus} (CD_3)(CH_3)$$
 (d)  $H_2C=CH-CD_3$   
24. Amongst the following amino acids, the (R)-enantiomer is represented by  
(a)  $NH_2 \stackrel{CH_3}{\longrightarrow} COOH$  (b)  $\stackrel{OOH}{\longrightarrow} OH_2$  (c)  $CH_3 \stackrel{NH_2}{\longrightarrow} (d) CH_3 \stackrel{COOH}{\longrightarrow} OH_2$   
25. Arrange the following halides in the decreasing order of  $S_N I$  reactivity  
 $CH_3CH_2CH_2CI$   $H_2C=CHCH(CI)CH_3$   $CH_3CH_2CH(CI)CH_3$   
(j) (II) (III)  
(a) 1 > 1I > 1II (b) II > 1 > III (c) II > II > 1 (d) III > II > 1  
26. The volume of 1N KMnO<sub>4</sub> required to reach equivalence point in the titration with  
0.01 mole of ferrous oxalate dissolved in dilute  $H_2SO_4$  is  
(a)  $3 \text{ cm}^3$  (b)  $30 \text{ cm}^3$  (c)  $10 \text{ cm}^3$  (d)  $20 \text{ cm}^3$   
(a)  $2 \text{ (b)} 3 \text{ cm}^3$  (c)  $10 \text{ cm}^3$  (d)  $20 \text{ cm}^3$   
27. The number of possible geometrical isomers for octahedral  $Co(ox)(PMe_3)_2NH_3CI$   
complex is  
(a)  $2 \text{ (b)} 3 \text{ cm}^3$  (c)  $10 \text{ cm}^3$  (d)  $20 \text{ cm}^3$   
28. The paramagnetic species among the following is:  
(a)  $B_2$  (b)  $C_2$  (c)  $O_2^2$  (d) CO  
29. The purple colour of iodine vapours is due to  
(a) d-d transition (b)  $\pi - \sigma^*$  transition  
(c) charge transfer transition (d)  $\pi^* + \sigma^*$  transition  
(d)  $CO_4^2^-$  (b)  $VO_4^3^-$  (c)  $FOQ_4^2^-$  (d)  $MnQ_4^2^-$   
31. <sup>19</sup>F-NMR spectrum of meriodional isomer of octahedral RhCl\_3F\_3 complex, assuming  
 $J_{Rh-F} > J_{F-T}$ , will show  $[^{108}Rh(I = \frac{1}{2}); ^{19}F(I = \frac{1}{2})]$   
(a) one doublet (b) two doublets and one triplet  
(c) two doublets and two triplets (d) one singlet and two triplets  
32. Which one of the following will show closo structure?  
(a)  $B_5H_9$  (b)  $[B_{12}H_{12}]^2^-$  (c)  $B_4H_{10}$  (d)  $B_5H_{11}$   
33. The correct order of energy level of d-orbitals in ferrocene is:  
(a)  $d_x^2-y^2, d_{xy} < d_{xz} < d_{xz} < d_{xz} < d_{xz}-y^2, d_{xy} < d_{xz} < d_{xz} < d_{xz} < d_{xz}-y^2, d_{xy} < d_{zz}$ 

 $\bigoplus$ 



34. Two moles of a monoatomic perfect gas initially at 4.0 bar and 47 °C undergoes reversible expansion in an insulated container. The temperature at which the pressure reduces to 3.0 bar is (a) 200 K (b) 285 K (c) 310 K (d) 320 K

35. The mean ionic activity coefficient of 0.0005 mol kg<sup>-1</sup> CaCl<sub>2</sub> in water at 25 °C is: (a) 0.98 (b) 0.67 (c) 0.81 (d) 0.91

For the cell: Cd(Hg) | CdSO<sub>4</sub>.  $\left(\frac{8}{3}\right)$  H<sub>2</sub>O (s) | CdSO<sub>4</sub> (aq, satd.) | Hg<sub>2</sub>SO<sub>4</sub>(s) | Hg 36.

The temperature dependence of emf in Volts is given by

 $B = 1.0185 + 4.05 \times 10^{-5} (T - 293) + 9.5 \times 10^{-7} (T - 293)^2$ 

The change in entropy at 25 °C for the cell reaction is

(b) 9.65  $JK^{-1}mol^{-1}$  (c)  $8.3 \times 10^{-4} JK^{-1}mol^{-1}$ (a)  $-253 \text{ K}^{-1} \text{mol}^{-1}$ (d) zero

- 37. Two separate bulbs contain ideal gases A and B respectively. The density of gas-A is twice that of gas-B and molecular weight of gas-A is half of that of gas-B. The ratio of pressure of gas-A to that of gas-B is (b) 6 (c) 4(a) 3 (d) 1
- 38. Choose the correct criterion of spontaneity in terms of the properties of the system alone.

(a) 
$$(dS)_{U,V} > 0$$
 (b)  $(dS)_{T,P} > 0$  27 00 (c)  $(dS)_{H,P} < 0$  (d)  $(dG)_{T,V} < 0$ 

- 39. Compared to  $C_2H_6$ , the value of Vander Waal's constants 'a' and 'b' for He will be (b) 'a' will be larger but 'b' will be smaller (a) both will be smaller (c) 'b' will be larger but 'a' will be smaller (d) both will be larger
- The number of hyperfine components observed in the electronic transition  ${}^{2}p_{1/2} \rightarrow$ **40.** 
  - ${}^{2}S_{1/2}$  of an atom with nuclear spin  $\frac{1}{2}$  is
    - (a) 3 (b) 4 (c) 6 (d) 5
- 41. Given that,

$$\rho_{n,t,m}(r,\theta,\varphi) = R_{nt}(r)Y_{tm}(\theta,\varphi); R_{20}(r) \propto (2-r/a_0)e^{-r/a_0} \ , Y_{0,0}(\theta,\varphi) = 1/\sqrt{4\pi}.$$

The position of radial node in the 2s orbital is at

(a)  $r = a_0$ (b)  $2r = 2a_0$ (c)  $r = a_0/2$ (d)  $r = a_0/4$ 

42. Consider the following reaction and use the data given below

 $N_2(g) + 3H_2(g) \rightarrow 2NH_3(g)$  $\Delta H^0$  (25 °C) = -92.2 kJ  $H_2(g)$ 

Substance  $N_2(g)$ 

 $\oplus$ 

 $NH_3(g)$ 



**C**<sub>n</sub>/(**JK**<sup>-1</sup>**mol**<sup>-1</sup>) 29.1 28.8 35.1

5

Assuming  $C_p$  to be independent of temperature, the reaction at 100 °C compared to that at 25 °C will be

(a) Endothermic (b) Less exothermic (c) More exothermic (d) Having  $\Delta H^0 = 0$ 

43. The reaction of 2-methylfuran with DMF–POCl<sub>3</sub> would give



44. The major product formed during the hydroboration-oxidation of 1-methyl cyclopentene is



- 45. Which one of the following carbonyl compounds will give a fragment ion at m/z = 58 in their mass spectra?
  - (a)  $C_2H_5CH(CH_3)CHO$  (b)  $CH_3CH_2CH_2CH_2CHO$
  - (c)  $CH_3CH_2CH_2COCD_3$   $UDY_{AM-TN-27-00}$  (d)  $(CH_3)_2CHCH_2CHO$
- 46. The decreasing order of reactivity of the following towards OH<sup>-</sup> ions is
  - (I) meta-nitrobromobenzene (II) 2,4,6-trinitrobromobenzene
  - (III) para-nitrobromobenzene (IV) 2, 4-dinitrobromobenzene

(a) I > II > III > IV (b) II > IV > III > I (c) IV > II > III > I (d) II > IV > I > III

47. Identify the isotactic polypropylene from the following

(a) 
$$H \xrightarrow{Me} H Me Me H$$
  
(c)  $H \xrightarrow{Me} H Me H$   
(c)  $H \xrightarrow{Me} H Me H$   
(c)  $H \xrightarrow{Me} H Me$ 

#### 48. Which one of the following compounds will form an osazone derivative?

- (a) CH<sub>3</sub>CH<sub>2</sub>COCH<sub>2</sub>OH
- (c) CH<sub>3</sub>CH<sub>2</sub>CHOHCH<sub>2</sub>OH
- (b) CH<sub>3</sub>COCH<sub>2</sub>CH<sub>2</sub>OH (d) CH<sub>3</sub>CH<sub>2</sub>COCH<sub>2</sub>OCH<sub>3</sub>
- 49. buta-1,3-diene on heating with maleic anhydride would give





50. The product obtained during the following photochemical reaction is



- 51. There are two isomers of Pt(NH<sub>3</sub>)<sub>2</sub> Cl<sub>2</sub>, X and Y. When X is reacted with thiourea(tu). [Pt(tu)<sub>4</sub>]<sup>2+</sup> is formed while Y on reaction with thiourea yields [Pt(NH<sub>3</sub>)<sub>2</sub>(tu)<sub>2</sub>]. Identify X and Y and explain the reaction.
- 52. Ligand substitution reaction on metal clusters are often found to occur by associative mechanism by breaking of a M-M bond and thereby providing an open coordination site for the incoming ligand. Which one of the two clusters,  $Co_4(CO)_{12}$  or  $Ir_4(CO)_{12}$  is expected to undergo faster exchange with <sup>13</sup>CO? Suggest an explanation.
- 53. Using crystal-field theory, account for the fact that in square pyramidal  $[Ni(CN)_5]^{3-}$  ion, the axial Ni-C bond (217 pm) is longer than Ni-C basal bonds (187 pm), while in trigonal bipyramidal  $[CuCl_5]^{3-}$  ion, the axial Cu-Cl bonds (229 pm) are shorter than the Cu-Cl equatorial ones (239 pm).
- 54. Between complexes [Co(NH<sub>3</sub>)<sub>5</sub>Cl]<sup>2+</sup> and [Co(NH<sub>3</sub>)<sub>5</sub>Br]<sup>2+</sup> which one should have a lower energy charge-transfer band and why?
- 55. Write the Russell-Saunders terms of the configuration p<sup>1</sup>d<sup>1</sup>and identify the ground term. Indicate population of electrons in different d-orbitals corresponding to <sup>3</sup>T<sub>1g</sub>(F), <sup>3</sup>T<sub>2g</sub>(F) and <sup>3</sup>A<sub>2g</sub>(F) states for d<sup>2</sup> configuration in octahedral symmetry.
- 56. Propose the structure of compounds A, B and C satisfying EAN rule in the following reactions:



57. Explain why 16 or 14 electron configurations are favoured over 18-electron configurations for the elements at the end of the transition series?

 $\oplus$ 



- 58. Calculate spin-orbit coupling parameter ( $\lambda$ ), for an octahedral nickel (II) complex exhibiting spin allowed d-d bands at 10,750 cm<sup>-1</sup>, 17,500 cm<sup>-1</sup> and 28,200 cm<sup>-1</sup> respectively. The experimentally determined magnetic moment is 3.2 BM.
- 59. Explain why pKa of  $(CH_3)_3SiOH$  (pKa  $\approx 11$ ) is lower than that of  $(CH_3)_3COH$  (pKa  $\approx 16$ )
- 60. Define capacity factor. What is the effect of large capacity factor on the separation of analytes? How can capacity factors be optimized in gas chromatography?
- 61. What prevents simple iron porphyrins from functioning as  $O_2$  carriers like haemoglobin?
- 62. Identify all the symmetry operations for HCHO with rotation axis as the z-axis and plane of the molecule being the yz plane. will the transition from an  $a_1$  to  $b_1$  orbital be allowed in HCHO? What will be the polarization of the corresponding  $b_1 \rightarrow a_1$  emission?
- 63. Using quantization condition for de Broglie wavelength on a ring, derive the expression for rotational energy levels of a rigid homonuclear diatomic rotor of bond length 2r.
- 64. Given  $\varphi_n(x) = (2)^{1/2} \sin(n\pi x)$ , show that eigen functions  $\varphi_1(x)$  and  $\varphi_2(x)$  of a particle in a one-dimensional box of length 1 are orthogonal.
- 65. The emf of the cell

Ag(s) | AgCl (satd), KCl (0.05 mol dm<sup>-3</sup>) | AgNO<sub>3</sub> (0.1 mol dm<sup>-3</sup>) | Ag(s) is 0.431 v at 298.15 K. The mean activity coefficient of KCl is 0.817 and that of AgNO<sub>3</sub> is 0.723. Calculate the solubility product of AgCl at 25 °C.

66. A paramagnetic substance  $(A_2B_3.5H_2O)$  initially at T = 0.30 K was magnetized by application of strong magnetic field while the sample was surrounded by helium gas in contact with a cold reservoir. Subsequently, helium gas was pumped away and the magnetic field was slowly reduced to zero. Calculate the change in temperature of the sample using the data given below:

Unma	gnetized Sample	Magnetized Sample			
T/K	$S/(JK^{-1}mol^{-1})$	T/K	S/(JK <sup>-1</sup> mol <sup>-1</sup> )		
0.30	0.40	0.30	0.19		
0.25	0.32	0.25	0.15		





0.18	0.19	0.20	0.12
0.15	0.14	0.15	0.10

- 67. When 2 moles of liquid-A and 4 moles of liquid-B are mixed, experimental measurements give entropy of mixing as 42 JK<sup>-1</sup>. Show whether the solution AB thus formed is ideal or not.
- 68. The LCAOs :  $\phi_1 = (\mathbf{1}_{s_A} + \mathbf{1}_{s_B})$  and  $\phi_2 = (\mathbf{1}_{s_A} \mathbf{1}_{s_B})$  approximate the lowest  $\sigma$  and  $\sigma^*$  orbitals of  $H_2^+$  respectively. Show that the  $\sigma$  LCAO is of g-type and  $\sigma^*$  of u type. [ $\mathbf{1}_{s_A}$  and  $\mathbf{1}_{s_B}$  are the  $\mathbf{1}_s$  orbitals centered on  $H_A$  and  $H_B$  of  $(H_A - H_B)^+$ ]
- 69. The fundamental vibrational frequency of HCl is 2885 cm<sup>-1</sup>. Assuming that HCl and DCl may be treated as Simple Harmonic Oscillator, calculate the fundamental frequency of DCl.
- 70. The conversion of A to B and C goes through the following mechanism

$$2A \xrightarrow{k_1} 1 \xrightarrow{k_2} B + C$$

Show that the equilibrium constant (K) of the overall reaction is,  $K = \frac{k_1 k_3}{k_2 k_4}$ 

- 71. The rate of the acid catalyzed hydrolysis of ethylacetate in HCl solution obeys the following rate law, Rate = -d[ester]/dt = k[ester][HCl] where k = 0.1 mol<sup>-1</sup>dm<sup>-3</sup>h<sup>-1</sup>. Neglecting any back reaction, calculate the time required for half the ester to be hydrolyzed if the initial concentration of ester and HCl are 0.02 mol dm<sup>-3</sup> and 0.01 mol dm<sup>-3</sup> respectively.
- 72. Using the data given below, calculate the equilibrium constant and enthalpy of the following reaction at 25 °C.

Substance	$\Delta G_F^0 / (kJmol^{-1})$	$\Delta S^0 / (JK^{-1}mol^{-1})$		
<b>NO</b> <sub>2</sub> (g)	51.31	240.06		
N <sub>2</sub> O <sub>4</sub> (g)	97.89	304.29		

 $2NO_2(g) \longrightarrow N_2O_4(g)$ 

73. When 1.0 mol of  $CH_4$  (g) is oxidized to carbon dioxide and water according to the reaction,  $CH_4$  (g) +  $2O_2$ (g)  $\rightarrow CO_2$  (g) +  $2H_2O$  (*l*). The corresponding thermodynamic parameters are:  $\Delta H^0 = -890$  kJ mol<sup>-1</sup>, and  $\Delta S^0 = -140.3$ 





JK<sup>-1</sup>mol<sup>-1</sup>. Assuming ideal gas behaviour, calculate the amount of energy that can be extracted as work at 25 °C.

74. Acid catalyzed dehydration of a tertiary alcohol-A (C<sub>6</sub>H<sub>14</sub>O) gives one major compound-B, and one minor compound-C both having molecular formula C<sub>6</sub>H<sub>12</sub>. Spectroscopic data of these compounds are as follows:

**Compound B :** 

IR :  $1660 \text{ cm}^{-1}$ ,  $3080 \text{ cm}^{-1}$ <sup>1</sup>H NMR  $\delta$  : 0.91(t, J = 7Hz, 3H), 1.60 (s, 3H), 1.70 (s, 3H), 1.98 (quin, J = 7 Hz, 2H), 5.08 (t, J = 7 Hz, 1H)

**Compound C :** 

IR :  $1640 \text{ cm}^{-1}$ ,  $3090 \text{ cm}^{-1}$ 

<sup>1</sup>H NMR  $\delta$ : 0.92 (t, J = 7 Hz, 3H), 1.40 (sextet, J = 7 Hz, 2H), 1.74 (s, 3H), 2.02 (t, J = 7 Hz, 2H), 4.78 (s, 2H)

Deduce the structure of A, B and C.

75. Suggest a plausible mechanism for the following reaction



76. Propose a mechanism for the photochemical reaction given below.



- 77. Draw  $\pi$  orbitals of buta-1,3-diene and ethylene, and identify their HOMO and LUMO.
- 78. Using "frontier orbital concept", explain why the  $\pi^{4s} + \pi^{2s}$  cycloaddition given below is photochemically not allowed?



79. Suggest a synthetic route to the hydroxy-ketone-C using A and B starting materials.



Tiruchirappalli – 620 024

ajchemacademy@gmail.com

 $\square$ 



80. The reaction of dimethyl fulvene-D with PhLi readily gives the anion-E. However, the analogous compound-F does not react with PhLi to give the corresponding anion-G. Explain.



- 81. Outline a synthesis of para-nitropropylbenzene from benzene.
- 82. Predict the products in the following reactions.

$$A \xrightarrow{Ph_3P=CH_2} B$$

83. Write the structure of the products X, Y and Z in the following sequence of reactions.

Stal -3

- 84. **D-Glucose** and **D-fructose** interconvert into each other in aqueous alkaline solution. Suggest a mechanism for the interconversion.
- 85. Propose a mechanism of the following reaction.



86. Suggest a route for synthesis of the enone 'B' from 'A'



# Answer Key

Q.No	Ans	Q.No	Ans	Q.No	Ans	Q.No	Ans
1.	d	16.	С	31.	С	46.	b
2.	С	17.	С	32.	b	47.	С

Tiruchirappalli – 620 024

www.csircoaching.com

ajchemacademy@gmail.com





3.	b	18.	d	33.	а	48.	а
4.	b	19.	b	34.	b	49.	b
5.	С	20.	а	35.	d	50.	С
6.	d	21.	d	36.	b		
7.	а	22.	d	37.	С		
8.	а	23.	d	38.	а		
9.	С	24.	b	39.	а		
10.	d	25.	С	40.	С		
11.	С	26.	С	41.	b		
12.	b	27.	С	42.	С		
13.	b	28.	а	43.	b		
14.	d	29.	d	44.	d		
15.	b	30.	С	45.	а		

#### ATEGIAM

© No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.

© Inspite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So, we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.

© இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.

© இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

 $\square$ 

