

GATE – 2004 – Chemistry



www.csircoaching.com

- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

Features

- | | |
|-------------------------------|----------------------------------|
| ➤ 300 ++ Live Classes | ➤ A Well-Defined Curriculum |
| ➤ 200 ++ Concept Wise Tests | ➤ A Strong Subject Foundation |
| ➤ 50 ++ Chapter Wise Tests | ➤ A Refined Learning Methodology |
| ➤ 50 ++ Model Tests | ➤ Updated Study materials |
| ➤ 2000 ++ Problem Discussions | ➤ Freshers Can easily understand |
| ➤ Recorded Videos | ➤ Question banks |



















Q.1 – Q.30 Multiple Choice Question (MCQ), carry ONE mark each (for each wrong answer: – 1/3).

- In units of $\frac{h^2}{8ml^2}$, the energy difference between levels corresponding to 3 and 2 node eigenfunctions for a particle of mass m in a one dimensional box of length ℓ is
(a) 1 (b) 3 (c) 5 (d) 7
- On the basis of LCAO-MO theory, the magnetic characteristics of N_2 and N_2^+ are
(a) N_2 is paramagnetic and N_2^+ is diamagnetic (b) both are paramagnetic
(c) N_2 is diamagnetic and N_2^+ is paramagnetic (d) both are diamagnetic
- The v_{rms} of a gas at 300 K is $30 R^{1/2}$. The molar mass of the gas, in $Kg\ mol^{-1}$, is
(a) 1.0 (b) 1.0×10^{-1} (c) 1.0×10^{-2} (d) 1.0×10^{-3}
- The coefficient of performance of a perfect refrigerator working reversibly between the temperature T_c and T_h is given by
(a) $\frac{T_c - T_h}{T_c}$ (b) $\frac{T_h - T_c}{T_c}$ (c) $\frac{T_c}{T_h - T_c}$ (d) $\frac{T_h}{T_h - T_c}$
- At a given temperature and pressure, the phase diagram of a three component system shows a binodal curve. If the two components are chloroform and water, the third component, among the choices given below, is
(a) benzene (b) acetic acid (c) toluene (d) carbon tetrachloride
- A certain reaction proceeds in a sequence of three elementary steps with the rate constants k_1 , k_2 and k_3 . If the observed rate constant (k_{obs}) of the reaction is expressed as $k_{obs} = k_3(k_1/k_2)^{1/2}$, the observed activation energy (E_{obs}) of the reaction is
(a) $\frac{1}{2} \left[\frac{E_1}{E_2} \right] + E_3$ (b) $\frac{E_3 + E_1}{E_2}$ (c) $E_3 \left[\frac{E_1}{E_2} \right]^{1/2}$ (d) $E_3 + \frac{1}{2} (E_1 - E_2)$
- Which one of the following is an example of a maximum boiling azeotrope?
(a) $H_2O - HCl$ (b) $H_2O - C_2H_6OH$ (c) $CHCl_3 - CH_3OH$ (d) $CCl_4 - CH_3OH$
- For the reaction, $A + B \leftrightarrow X^\ddagger \rightarrow P$, $E_a = 20.0\ kJ\ mol^{-1}$ at 300K. The enthalpy change for the formation of the activated complex from the reactants in $kJ\ mol^{-1}$ is
(a) 12 (b) 15 (c) 23 (d) 25
- In an osmotic pressure measurement, a plot of height of solution (h) of density (ρ) versus concentration ($g\ L^{-1}$) was made at a temperature T . The slope of the plot will be equal to (where g , given in the choices below, is the acceleration of free fall)
(a) $\frac{\rho RT}{gM}$ (b) $\frac{gRT}{\rho M}$ (c) $\frac{RT}{\rho gM}$ (d) $\frac{gRM}{\rho T}$




10. If 0.001 M of a substance quenches the efficiency of fluorescence by 20%, the value of Stem-Volmer constant in M^{-1} is
 (a) 100 (b) 150 (c) 200 (d) 250
11. Which one of the following is NOT a photodetector?
 (a) Bolometer (b) Charge-transfer device
 (c) Photomultiplier tube (d) Silicon diode
12. The nature of excitation signal used for cyclic voltammetry is
 (a) linear scan (b) differential pulse (c) triangular (d) square wave
13. The structure of SF_4 is
 (a) octahedral (b) tetrahedral (c) trigonal bipyramidal (d) square planar
14. The number of metal-metal bonds present in $Ir_4(CO)_{12}$ are
 (a) 4 (b) 5 (c) 6 (d) 8
15. The zero magnetic moment of octahedral $K_2[NiF_6]$ is due to
 (a) low spin d^6 Ni(IV) complex (b) low spin d^8 Ni(II) complex
 (c) high spin d^8 Ni(II) complex (d) high spin d^6 Ni(IV) complex
16. The number of hyperfine split lines observed in ESR spectrum of methyl radical is
 (a) 1 (b) 4 (c) 6 (d) 8
17. The absorption of $[Co(NH_3)_6]^{2+}$ is:
 (a) stronger than that of $[Co(NH_3)_5Cl]^{2+}$
 (b) stronger than that of $[MnCl_4]^{2-}$
 (c) weaker than that of $[MnCl_4]^{2-}$ but stronger than that of $[Co(NH_3)_5Cl]^{2+}$
 (d) weaker than those of both $[MnCl_4]^{2-}$ and $[Co(NH_3)_5Cl]^{2+}$
18. Which one of the following statements about ferrocene is FALSE?
 (a) It obeys the 18-electron rule (b) It is diamagnetic
 (c) It is an orange solid (d) It resists electrophilic substitution
19. The bond angle of Cl_2O is
 (a) smaller than that of F_2O (b) greater than that of H_2O
 (c) smaller than that of H_2O (d) same as that of F_2O
20. The half-wave potential for a reversible reduction of a metal ion in polarography is independent of
 (a) Concentration of the supporting electrolyte
 (b) Concentration of the electroactive species
 (c) Concentration of the complexing agent




- (d) Temperature of the solution
21. The major product formed on **nitration of N,N-dimethylaniline with conc. $\text{H}_2\text{SO}_4 - \text{HNO}_3$ mixture is**
- (a)  (b)  (c)  (d) 
22. **Reaction of phenylacetylene with sodamide in liquid ammonia generates**
- (a)  (b)  (c)  (d) 
23. **Proton decoupled ^{13}C -NMR spectrum of a bicyclooctane (C_8H_{14}) exhibits only two signals. The Structure of the compound is:**
- (a)  (b)  (c)  (d) 
24. **Cyclohexyl benzyl ether when reacted with hydrogen in the presence of 10% palladium on charcoal generates a mixture of**
- (a) cyclohexanol and benzyl alcohol (b) cyclohexane and benzyl alcohol
(c) cyclohexanol and toluene (d) cyclohexane and toluene
25. **In electrophilic aromatic substitution reactions, nitro group is meta-directing, because the nitro group**
- (a) increase electron density at meta-position
(b) increase electron density at ortho-and para-positions
(c) decreases electron density at meta-position
(d) decreases electron density at ortho-and para-positions
26. **Among the resonance forms given below, the one which contributes most to the stability of azulene is**
- (a)  (b)  (c)  (d) 
27. **The configurations at the two asymmetric centres (C-1 and C-6) in the bicyclo[4.4.0]decane, given below are**





- (a) 1R, 6R (b) 1R, 6S (c) 1S, 6S (d) 1S, 6R
28. The **reactive intermediate** involved in the conversion of **phenol** to **salicylaldehyde** using **chloroform** and **sodium hydroxide** is
- (a) $\text{Cl}_2\text{C}:$ (b) Cl_2CH^+ (c) Cl_2CH^- (d) Cl_2CH^+
29. Conversion of **Ph-NH₂** into **Ph-CN** can be accomplished by
- (a) reaction with sodium cyanide in the presence of nickel catalyst
 (b) reaction with chloroform and sodium hydroxide
 (c) diazotisation followed by reaction with CuCN
 (d) reaction with ethyl formate followed by thermolysis
30. The **vicinal coupling constant (J)** expected for the protons **H_P** and **H_Q** in the compound given below will be in the range



- (a) 0 - 2 Hz (b) 4 - 6 Hz (c) 8 - 10 Hz (d) 12 - 15 Hz

Q.31 – Q.90 Multiple Choice Question (MCQ), carry TWO marks each (for each wrong answer: – 2/3).

31. For one mole of an ideal gas, $\left(\frac{\partial P}{\partial T}\right)_V \left(\frac{\partial V}{\partial T}\right)_P \left(\frac{\partial V}{\partial P}\right)_T =$
- (a) -1 (b) $-\frac{R^2}{P^2}$ (c) +1 (d) $\frac{R^2}{P^2}$
32. Neglecting the mass of **hydrogen (1.0 amu)** and **deuterium (2.0 amu)** with respect to that of **iodine (127 amu)**, the ratio between **fundamental vibrational frequencies of HI and DI** is:
- (a) $\frac{1}{2}$ (b) 2 (c) $\frac{1}{\sqrt{2}}$ (d) $\sqrt{2}$
33. The population of **Jth rotational level N_J** is given by $N_J = N_0(2J + 1)e^{[J(J+1)B]/KT}$. The **J value of rotational level with maximum population (J_{max})** is given by
- (a) $\frac{(2kT/B)-1}{\sqrt{2}}$ (b) $\frac{\sqrt{2kT/B}-1}{2}$ (c) $\frac{kT}{B}$ (d) $\frac{B}{kT}$
34. The **fugacity coefficient (φ)** is given by $\ln \phi = \int_0^P \left(\frac{z-1}{p}\right) dp$ where **z** is the



- compressibility factor**, and p the pressure. The fugacity of a gas governed by the gas law $p(V_m - b) = RT$ is
- (a) $p \ln (V_m/RT)$ (b) $pe^{b/RT}$ (c) $pe^{-bp/RT}$ (d) $pe^{bp/RT}$
35. The **number and symmetry type of normal modes of vibration of H_2O** are
- (a) 3 and $2A_1 + B_2$ (b) 3 and $2A_1 + A_2$
 (c) 3 and $2A_1 + B_1$ (d) 4 and $3A_1 + B_2$
36. The gaseous reaction $2A + B \rightarrow C$, with partial pressures of $p_A = 0.1 \text{ atm}$; $p_B = 0.001 \text{ atm}$ and $p_C = 1.0 \text{ atm}$, proceeds to the left at 298 K. The **equilibrium constant, K_p** for the above reaction is
- (a) 1.0×10^4 (b) 1.0×10^5 (c) 1.0×10^6 (d) 1.0×10^7
37. The **change in entropy** when one mole of an ideal gas is compressed to one-fourth of its initial volume and simultaneously **heated to twice its initial temperature** is
- (a) $(C_v - R) \ln 4$ (b) $(C_v - 2R) \ln 2$ (c) $(C_v - 2R) \ln 4$ (d) $(C_v + 2R) \ln 2$
38. For the reaction, $A_{(s)} \leftrightarrow B_{(l)} + 2C_{(g)}$, $\Delta G^0 \text{ (in Joules)} = 90800 - 100T$. The **partial pressure of $C_{(g)}$ at 600 K in Torr** is
- (a) 15 (b) 22 (c) 35 (d) 46
39. Match the following:
- | | | | | | |
|--|------------------|-----|-----|-----|----|
| P. $\left(\frac{\partial U}{\partial S}\right)_V$ | (i) V | | | | |
| Q. $\left(\frac{\partial U}{\partial V}\right)_S$ | (ii) $-S$ | (a) | iii | iv | i |
| R. $\left(\frac{\partial G}{\partial P}\right)_T$ | (iii) T | (b) | iii | i | ii |
| S. $\left(\frac{\partial G}{\partial T}\right)_P$ | (iv) $-P$ | (c) | i | iii | iv |
| | | (d) | iv | iii | i |
40. Match the following :
- | | |
|-----------------------------------|----------------------------------|
| P. $4n + 2$ rule | I. Woodward-Hoffmann rule |
| Q. single valued | II. Bound system |
| R. $< p_x \geq 0$ | III. Hurtle-Fock Theory |
| S. photochemically allowed | IV. Huckel theory |
| | V. Wave function |
| | VI. unbound system |

P Q R S

P Q R S

- (a) I ; III ; IV ; VI (b) IV ; V ; II ; VI
(c) II ; VI ; III ; I (d) IV ; V ; II ; I
41. The **solubility product of silver sulphate** at 298 K is 1.0×10^{-5} . If the **standard reduction potential** of the halfcell $\text{Ag}^+ + e \rightarrow \text{Ag}$ is 0.80 V, the **standard reduction potential** of the half-cell $\text{Ag}_2\text{SO}_4 + 2e \rightarrow 2\text{Ag} + \text{SO}_4^{2-}$ is:
(a) 0.15 V (b) 0.22 V (c) 0.65 V (d) 0.95 V
42. The **criterion for spontaneous change** in terms of the state functions is:
(a) $dU_{S,V} \geq 0$ (b) $dA_{T,V} \geq 0$ (c) $dS_{U,V} \geq 0$ (d) $dG_{T,V} \leq 0$
43. One mole of an ideal gas ($C_V = 1.5R$) at a temperature 500 K is compressed from 1.0 atm to 2.0 atm by a reversible isothermal path. Subsequently, it is expanded back to 1.0 atm by a **reversible adiabatic path**. The volume of the final state in litre is:
(a) 15.6 (b) 20.5 (c) 31.1 (d) 41.0
44. The **vapour pressures of the pure components 'P' and 'Q'** are 700 Torr and 500 Torr, respectively. When the two phases are in equilibrium at 1.0 atm, the **mole fraction of 'P' in the liquid phase is 0.6 and in the vapour phase 0.4**. The **activity coefficient of component-P in the solution on the basis of Raoult's law** is
(a) 0.60 (b) 0.72 (c) 0.92 (d) 1.01
45. The **concentration of oxygen in water in mg L^{-1}** . If the **Henry's law constant for oxygen at 298 K is 2.80×10^7 Torr**, the **partial pressure of oxygen in the atmosphere in Torr** is,
(a) 28 (b) 32 (c) 50 (d) 15.68
46. **Decomposition of ammonia on tungsten at 850 °C** has a rate constant value of 0.10 Torr s^{-1} . If the initial pressure of ammonia is 100 Torr, **the pressure of ammonia (in Torr) at $t = 200$ s** is
(a) 10 (b) 20 (c) 50 (d) 80
47. For the reaction of the type $\text{P} \xrightarrow{k_1} \text{Q} \xrightarrow{k_2} \text{R}$, given that $[\text{P}]_0 = 1.0\text{M}$; $k_1 = 1 \times 10^{-3} \text{s}^{-1}$ and $k_2 = 1 \times 10^{-4} \text{s}^{-1}$, the time at which the concentrations of **Q and R are 0.5966 M and 0.0355M, respectively**, is
(a) 500s (b) 750s (c) 1000s (d) 1500s
48. The spinels **CoFe_2O_4 and FeFe_2O_4** , respectively, are
(a) inverse and inverse (b) inverse and normal



- (c) normal and normal (d) normal and inverse
49. According to **Wade's rule**, the structures of $\text{B}_{10}\text{C}_2\text{H}_{12}$ and $[\text{B}_9\text{C}_2\text{H}_{11}]^{2-}$, respectively, are
 (a) closo and arachno (b) nido and closo (c) closo and nido (d) nido and arachno
50. The **overall charge** present on the **cyclic silicate anion** $[\text{Si}_6\text{O}_{18}]^{n-}$ is
 (a) 6 (b) 12 (c) 18 (d) 24
51. The **ground state term symbols** for **high spin** d^5s^1 and d^5 configurations, respectively, are
 (a) 3S and 6S (b) 6P and 3S (c) 7S and 6S (d) 7P and 6S
52. The **reagents required for the synthesis of cyclic phosphazene** $\text{N}_4\text{P}_4\text{Cl}_8$ are
 (a) PCl_5 and NH_3 (b) POCl_3 and NH_4Cl
 (c) POCl_3 and NH_3 (d) PCl_5 and NH_4Cl
53. The **isomerisms** that are possible in the **Co(III) complexes** $[\text{Co}(\text{NH}_3)_3(\text{NO}_2)_3]$ and $[\text{Co}(\text{NH}_3)_5\text{NO}_2]\text{Cl}_2$, respectively, are
 (a) co-ordination and position (b) optical and linkage
 (c) geometrical and linkage (d) optical and optical
54. The **perxenate ion** $[\text{XeO}_6]^{4-}$ can be prepared by
 (a) direct reaction of Xe with oxygen (b) reaction of XeF_6 with oxygen
 (c) hydrolysis of XeF_6 in acidic medium (d) hydrolysis of XeF_6 in basic medium
55. In **tetrahedral geometry**, which one of the following sets of **electronic configurations** will have **orbital contribution to the magnetic moment**?
 (a) d^3, d^4, d^8 and d^9 (b) d^1, d^6, d^7 and d^9 (c) d^3, d^4, d^7 and d^9 (d) d^1, d^3, d^4 and d^9
56. The most suitable route to prepare **trans**- $[\text{PtCl}_2(\text{NH}_3)(\text{PPh}_3)]$ is:
 (a) $[\text{PtCl}_4]^{2-}$ with PPh_3 followed by reaction with NH_3
 (b) $[\text{PtCl}_4]^{2-}$ with NH_3 followed by reaction with PPh_3
 (c) $[\text{Pt}(\text{NH}_3)_4]^{2+}$ with HCl followed by reaction with PPh_3
 (d) $[\text{Pt}(\text{NH}_3)_4]^{2+}$ with PPh_3 followed by reaction with HCl
57. A solution containing **5 ppm of KMnO_4 (M.W = 159)** has a **transmittance of 0.360** measured in a **1 cm cell at 500 nm**. The **molar absorptivity of KMnO_4 in $\text{L mol}^{-1} \text{cm}^{-1}$** is
 (a) 1.1×10^4 (b) 1.4×10^4 (c) 1.9×10^4 (d) 2.7×10^4



58. Match the following:

- | | |
|-----------------------------------|--------------------------------------|
| P. Coulometry | I. Dropping mercury electrode |
| Q. Ion selective electrode | II. Current efficiency |
| R. Polarography | III. Dead stop end point |
| S. Amperometry | IV. Membrane potential |
| | V. Conductometer |
| | VI. Actinometer. |

| | P | Q | R | S |
|-----|----|----|-----|-----|
| (a) | II | IV | I | III |
| (c) | VI | V | III | IV |

| | P | Q | R | S |
|-----|-----|----|-----|----|
| (b) | I | II | III | V |
| (d) | III | IV | I | VI |

59. Match the following:

- | | |
|-----------------------------------|---------------------------------|
| P. Ferritin | I. electron transport |
| Q. Vitamin B ₁₂ | II. Ionophore |
| R. Cytochromes | III. Oxygen transport |
| S. Valinomycin | IV. Nitrogen fixation |
| | V. Organometallic enzyme |
| | VI. Iron storage. |

| | P | Q | R | S |
|-----|-----|----|----|----|
| (a) | VI | IV | II | I |
| (c) | III | V | IV | VI |

| | P | Q | R | S |
|-----|----|-----|----|----|
| (b) | I | III | VI | IV |
| (d) | VI | V | I | II |

60. The number of absorption bands observed $[\text{FeF}_6]^{3-}$ and $[\text{CoF}_6]^{3-}$, respectively, are







- (a) 1 and 3 (b) 0 and 1 (c) 0 and 3 (d) 3 and 1

61. Regarding the catalytic cycle of hydrogenation of alkenes involving $[\text{RhCl}(\text{PPh}_3)_3]$ as the catalyst, the correct statements is:


- (a) Only 18-electron Rh complex is involved
 (b) 14-, 16- and 18-electron Rh complexes are involved
 (c) 14- and 16-electron Rh complexes are involved
 (d) 16- and 18-electron Rh complexes are involved

62. The IR stretching frequency (ν_{CO}) of P–S follows the order












- (a) $P > R > S > Q$ (b) $S > P > R > Q$ (c) $Q > S > P > R$ (d) $R > Q > P > S$
63. The structures of $N(CH_3)_3$ and $N(SiH_3)_3$ respectively, are
 (a) trigonal planar and pyramidal (b) pyramidal and trigonal planar
 (c) pyramidal and pyramidal (d) trigonal planar and trigonal planar
64. Which one of the following is **NOT** correct in chromatography?
 t_M = Retention time for a species that is not retained by the stationary phase
 t_R = Retention time for the analyte
 $(t_R)_n$ = Retention time for the component 'n'
 W_n = Width of the peak at its base for the component 'n'
 (a) Resolution = $\frac{(t_R)_2 - (t_R)_1}{2(W_1 + W_2)}$ (b) Capacity factor = $\frac{t_R - t_M}{t_M}$
 (c) Separation factor = $\frac{(t_R)_2 - t_M}{(t_R)_1 - t_M}$ (d) No. of theoretical plates = $16 \left(\frac{t_R}{W} \right)^2$
65. Thermal reaction of allyl phenyl ether generates a mixture of ortho-and para-allyl phenols. The para-allyl phenol is formed via
 (a) [3,5]-sigmatropic shift
 (b) first ortho-allyl phenol is formed, which then undergoes a [3,3]-sigmatropic shift
 (c) two consecutive [3,3]-sigmatropic shifts
 (d) dissociation to generate allyl cation, which then adds at para-position
66. Of the favour vicinal diols shown below, only three are cleaved by HIO_4 , the diol which is **NOT** cleaved HIO_4 is
 (a)  (b)  (c)  (d) 
67. With respect to the two reactions shown below, the correct statements about their stereochemical nature is $[LDA = LiN(iPr)_2]$
 (i)  (ii) 
 (a) The reactions are stereoselective, because P and Q are the same
 (b) The reactions are non-stereoselective, because P and Q are the same
 (c) The reactions are stereoselective, because P and Q are diastereomers
 (d) The reactions are enantioselective, because P and Q are enantiomers
68. For the reactions shown below, identify the correct statement with regard to the

products formed.




- (a) P and Q are identical, both are optically active
 (b) P and Q are positional isomers, P is racemic and Q is optically active
 (c) P and Q are positional isomers, P is optically active and Q is racemic
 (d) P and Q are positional isomers, both are optically active
69. In the reaction shown below, identify the correct combination of the **intermediate-P** and the **product-Q**.




- (a) P =  and Q = 
- (b) P =  and Q = 
- (c) P =  and Q = 
- (d) P =  and Q = 

70. In the two step reaction shown below, identify the **correct combination of products 'P' and 'Q'**.





71. On the basis of **Favorskii rearrangement** mechanism, the ratio of the **products P, Q and R** given below, will be, respectively.




- (a) 2 : 1 : 1 (b) 1 : 1 : 1 (c) 1 : 0 : 1 (d) 0 : 1 : 1
72. An organic compound having molecular formula $\text{C}_6\text{H}_{11}\text{BrO}_2$ exhibits the following peaks.


$^1\text{H-NMR}$: δ 4.1 (2H, q, $J = 7.5$ Hz), 4.0 (2H, t, $J = 7.5$ Hz), 1.5 - 2.2 (4H, m), 1.25 (3H, t, $J = 7.5$ Hz)

The structure of the compound is:




73. The **product-P** formed in the following three steps reaction is:





74. Identify the correct choice of reagents, among P, Q and R, for the transformation of norbornene into the epoxides I and II


P = $\text{H}_2\text{O}_2 - \text{AcOH}$; Q = $\text{H}_2\text{O}_2 - \text{NaOH}$; R = HOBr followed by aq. NaOH



- (a) P gives I and Q gives II (b) R gives I and P gives II
(c) Q gives I and R gives II (d) P gives I and R gives II
75. Reaction of ethyl acetoacetate with one equivalent of methylmagnesium bromide gives



76. For the aldotetroses I-IV, the combination of TRUE statements, among P-T, is:



P = I and II are diastereomers and II and III are enantiomers

Q = I and IV are mesomers and are optically inactive

R = I and III can be interconverted by a base catalysed isomerisation

S = Only I and IV are HIO_4 cleavable

T = I and III are D-sugars and II and IV are L-sugars

- (a) Q, R, T (b) P, R, T (c) Q, S, T (d) P, Q, S
77. Match the compounds P-S with their carbonyl stretching frequencies (cm^{-1}) I-VI in IR spectroscopy.

P. Acetone

I. 1870

P


Q

R


S

| | | | | | | |
|---------------------------|------------------|-----|-----|-----|-----|----|
| Q. ethyl acetate | II. 1800 | (a) | IV | III | I | VI |
| R. acetamide | III. 1740 | (b) | III | VI | V | II |
| S. acetyl chloride | IV. 1700 | (c) | IV | III | V | II |
| | V. 1660 | (d) | II | V | III | VI |
| | VI. 1600 | | | | | |




78. In the following three step transformation, identify the correct combination of product **P**, **Q** and **R**. [LDA = $\text{LiN}(\text{iPr})_2$].



79. The major product-**P** formed in the following photochemical reaction is:




80. Three molecular ionic states, **P-R**, are possible for the amino acid histidine. Identify the correct choice of pH values, respectively, for the observation of the ionic states **P-R**.

**P****Q****R**

- (a) P at pH 1 ; Q at pH 12 ; R at pH 7
 (b) P at pH 7 ; Q at pH 1 ; R at pH 12
 (c) P at pH 12 ; Q at pH 7 ; R at pH 1
 (d) P at pH 12 ; Q at pH 1 ; R at pH 7

81. In the reaction shown below, identify the correct combination of the **intermediate-P** and the **product-Q**.



Q 82 – 90 contain a Statement with a Reason and an Assertion. for each question, choose the correct answer from the given four choices.

82. **Statement** : Solid carbon dioxide is called as dry ice

Reason : CO_2 sublimates when kept in open atmosphere

Assertion : Triple point of CO_2 lies above one atmosphere

- (a) Both Reason and Assertion are correct
 (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong

- (d) Reason is wrong but Assertion is correct
83. **Statement** : Entropy of pure, perfectly crystalline substance is zero at absolute zero of temperature
- Reason** : At absolute zero, molecules can have only one orientation
- Assertion** : Statistical definition of entropy is given by the equation, $S = k \ln W$, where W is the probability of orientation
- (a) Both Reason and Assertion are correct
 (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong
 (d) Reason is wrong but Assertion is correct
84. **Statement** : Catalytic decomposition of ammonia on platinum takes place at 1000 °C
- Reason** : Ammonia is more strongly adsorbed than hydrogen on platinum
- Assertion** : The rate law for the decomposition of ammonia on platinum is given as, $\text{Rate} = k \frac{P_{\text{NH}_3}}{P_{\text{H}_2}}$
- (a) Both Reason and Assertion are correct
 (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong
 (d) Reason is wrong but Assertion is correct
85. **Statement** : $[\text{CoCl}_4]^{2-}$ is a regular tetrahedron but $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$ is a distorted
- Reason** : Unsymmetrical distribution of electrons in e_g orbital leads to distortion in $[\text{Cu}(\text{H}_2\text{O})_6]^{2+}$
- Assertion** : H_2O ligands interact differently with orbitals of unequal electron population. This leads to distortion in geometry
- (a) Both Reason and Assertion are correct
 (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong
 (d) Reason is wrong but Assertion is correct
86. **Statement** : Schottky and Frenkel defects are stoichiometric defect occurring in crystal lattices
- Reason** : Schottky defects are due to the absence of one positive and one negative ion and Frenkel defects are due to the presence of one



hole and one ion in an interstitial position

Assertion : The ratio of number of atoms of one kind to the number of atoms of the other kind does not correspond exactly to the ideal whole number ratio implied by the formula which leads to stoichiometric defects

- (a) Both Reason and Assertion are correct
- (b) Both reason and Assertion are wrong
- (c) Reason is correct and Assertion is wrong
- (d) Reason is wrong but Assertion is correct

87. **Statement :** Ga is below Al in Group IIIA, yet the atomic size of Ga is almost the same as that of Al

Reason : Lanthanide contraction

Assertion : Poor shielding of nuclear charge results in outer electrons being more firmly held by the nucleus

- (a) Both Reason and Assertion are correct
- (b) Both reason and Assertion are wrong
- (c) Reason is correct and Assertion is wrong
- (d) Reason is wrong but Assertion is correct

88. **Statement :** 5-Bromopyrimidine ($C_4H_3BrN_2$) exhibits two prominent peaks in the mass spectrum at m/z 158 and 160 in the ratio of 1:1

Reason : There are two basic centres in the molecule, which are protonated

Assertion : There are two isotopes of bromine, ^{79}Br and ^{81}Br , that occur in the ratio of 1:1

- (a) Both Reason and Assertion are correct
- (b) Both Reason and Assertion are wrong
- (c) Reason is correct and Assertion is wrong
- (d) Reason is wrong but Assertion is correct

89. **Statement :** Pyridine is more basic than pyrrole


Reason : The nitrogen in pyrrole carries a proton while the nitrogen in pyridine does not

Assertion : Nitrogens in trigonal geometry are generally more basic than the nitrogens in tetrahedral geometry

- (a) Both Reason and Assertion are correct



- (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong
 (d) Reason is wrong but Assertion is correct
90. **Statement** : Replacement of CH_3 with CF_3 decreases the rate decreases the rate of reaction-I, but increases the rate of reaction-II



Reason : Reaction-I proceeds through $\text{S}_{\text{N}}1$ mechanism and reaction-II proceeds through $\text{S}_{\text{N}}2$ mechanism

Assertion : Being an electron withdrawing group, CF_3 destabilizes the transition state in $\text{S}_{\text{N}}1$ reaction, but stabilizes the transition state in $\text{S}_{\text{N}}2$ reaction

- (a) Both Reason and Assertion are correct
 (b) Both reason and Assertion are wrong
 (c) Reason is correct and Assertion is wrong
 (d) Reason is wrong but Assertion is correct

Answer Key

| Q.No | Ans | | Q.No | Ans | | Q.No | Ans | | Q.No | Ans |
|------|-----|--|------|-----|--|------|-----|--|------|-----|
| 1. | d | | 26. | b | | 51. | c | | 76. | b |
| 2. | c | | 27. | b | | 52. | d | | 77. | c |
| 3. | d | | 28. | c | | 53. | c | | 78. | b |
| 4. | c | | 29. | c | | 54. | d | | 79. | c |
| 5. | c | | 30. | c | | 55. | a | | 80. | a |
| 6. | d | | 31. | b | | 56. | a | | 81. | b |
| 7. | a | | 32. | d | | 57. | b | | 82. | a |
| 8. | b | | 33. | b | | 58. | a | | 83. | a |

GATE – 2004 – CY

| | | | | | | | | | | |
|-----|---|--|-----|---|--|-----|---|--|-----|---|
| 9. | c | | 34. | d | | 59. | d | | 84. | a |
| 10. | d | | 35. | c | | 60. | b | | 85. | a |
| 11. | b | | 36. | b | | 61. | d | | 86. | c |
| 12. | b | | 37. | b | | 62. | d | | 87. | d |
| 13. | c | | 38. | b | | 63. | b | | 88. | d |
| 14. | c | | 39. | a | | 64. | a | | 89. | b |
| 15. | a | | 40. | d | | 65. | c | | 90. | a |
| 16. | b | | 41. | c | | 66. | a | | | |
| 17. | b | | 42. | c | | 67. | a | | | |
| 18. | d | | 43. | c | | 68. | d | | | |
| 19. | b | | 44. | b | | 69. | b | | | |
| 20. | d | | 45. | d | | 70. | c | | | |
| 21. | a | | 46. | b | | 71. | d | | | |
| 22. | b | | 47. | c | | 72. | c | | | |
| 23. | d | | 48. | a | | 73. | d | | | |
| 24. | c | | 49. | c | | 74. | d | | | |
| 25. | d | | 50. | b | | 75. | a | | | |

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © In spite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.

- © இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

