

JAM - Chemistry - 2006 (CY)



www.csircoaching.com

- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

Features

- > 300 ++ Live Classes
- > 200 ++ Concept Wise Tests
- > 50 ++ Chapter Wise Tests
- \succ 50 ++ Model Tests
- > 2000 ++ Problem Discussions
- Recorded Videos

- ➢ A Well-Defined Curriculum
- ➢ A Strong Subject Foundation
- ➢ A Refined Learning Methodology
- > Updated Study materials
- ➢ Freshers Can easily understand
- Question banks





JAM-2006-CY

<u>Attempt ALL the questions. Q.1 – Q.30 Multiple Choice Question</u> (MCQ), carry THREE marks each (for each wrong answer: –1).

1. After the following interchanges of groups in the Fischer projection of **2-bromobutane**, the configuration of (X) and (Y) will be



0

www.csircoaching.com

ajchemacademy@gmail.com



$$C_6H_6 + CO + HCl \xrightarrow{AlCl_3 / Cu_2Cl_2}$$

gives positive test with Fehling's solution. The product is

(a)
$$C_{6}H_{5}OH$$
 (b) $C_{6}H_{4}(CI)CHO$ (c) $C_{6}H_{4}(OH)CHO$ (d) $C_{6}H_{5}CHO$
6. The compound (X) in the reaction sequence is

$$\begin{array}{c} CH_{2}COONa & P_{2}S_{3} \\ CH_{2}COONa & heat \end{array} (X) & H_{2}, Raney Ni \\ CH_{3}CH_{2}CH_{2}CH_{2}CH_{3} \\ (a) & \swarrow \end{array} (b) & CH_{2}COOH \\ CH_{2}COOH & (c) & \swarrow \end{array} (d) & \swarrow \\ S \end{array}$$
7. The major product of the reaction is
(a) $(b) & (b) & (b) & (b) & (c) & (c)$

8. The increasing order of the acidity of the hydrogen marked in **bold italics** among the following is

н Ή Π III Ι (a) III < Π Ι (b) < I < III <Π (c) Ι < Π < III (d) Π < III < I The major product of the reaction is HBr

2

9.



ً⊘



19.	The optical absorption	on spectrum of [Ti(H ₂ O	$[b_{6}]^{3+}$ has its absorptio	n maximu			
(a) 1		(b) 3/2	(c) 3/4	(d) 4/3			
18. The average value of C-C bond order in graphite is							
	(a) 850	(b) 1600	(c) 200	(d) 400			

19. The optical absorption spectrum of [Ti(H₂O)₆]³⁺ has its absorption maximum at 20300 cm⁻¹. The magnitude of crystal field stabilization energy in cm⁻¹ is
(a) 8120 (b) 16240 (c) 24360 (d) 50750

- 20. In inorganic qualitative analysis, H2S in acidic medium will NOT precipitate(a) HgS(b) ZnS(c) CuS(d) CdS
- 21. The phase diagram of a pure substance is sketched below.



The number of degrees of freedom at points P1, P2 and P3, respectively, are

(a) 2, 1, 0 (b) 1, 2, 0 (c) 2, 0, 1 (d) 0, 2, 1 (d) 0, 2, 1

22. The solubility products (K_{sp}) for three salts MX, MY₂ and MZ₃ are 1×10^{-8} , 4×10^{-9} and 27×10^{-8} , respectively. The solubilities of these salts follow the order

(a)	MX	>	MY_2	>	MZ ₃	AUF	(b)	MZ_3	>	MY_2	>	MX
(c)	MZ_3	>	MX	>	MY ₂		(d)	MY_2	>	MX	>	MZ_3

- 23. The temperature (T) dependence of the equilibrium constant (K) of a chemical reaction is correctly described by the following statement:
 - (a) For an endothermic reaction, the slope of $\ln K \text{ vs } 1/T$ plot is positive
 - (b) For an exothermic reaction, K is proportional to T
 - (c) For an exothermic reaction, K at a higher temperature is lower than K at a lower temperature
 - (d) If ΔH is independent of temperature, the change in K with T is smaller at lower temperatures
- 24. When the concentration of K⁺ across a cell membrane drops from 0.01 M to 0.001 M, the potential difference across the membrane is

Tiruchirappalli – 620 024





JAM - 2006 - CY



5

The wavefunction for a particle (moving in a ring) is $(2\pi)^{-1/2} \exp(2i\phi)$, where 27. ϕ is the polar angle. The probability of finding the particle in a small interval d ϕ when the value of $\phi = \pi / 2$ is

- (b) $d\phi/2\pi$ (c) $d\phi \exp(i\pi)$ (a) $d\phi$ (d) $d\phi \exp(i\pi)/2\pi$
- An electric current of 0.965 ampere is passed for 2000 seconds through a 28. solution containing $[Cu(CH_3CN)_4]^+$ and metallic copper is deposited at the cathode. The amount of Cu deposited is
 - (c) 0.02 mol (b) 0.01 mol (a) 0.005 mol (d) 0.04 mol
- 29. The Maxwell-Boltzmann distribution for molecular speeds is shown in the following figure.



In the figure, H is the height of the peak, L is the location of the maximum and W is the width at half height. As the temperature is decreased,

- (a) H increases, L decreases and W increases
- (b) H increases, L decreases and W decreases
- (c) H decreases, L decreases and W increases
- (d) H decreases, L decreases and W decreases

(a) 0.0 V

25.

26.

(a) N_2





- **30.** A system undergoes two cyclic processes 1 and 2. Process 1 is reversible and process 2 is irreversible. The correct statement relating to the two processes is
 - (a) ΔS (for process 1) = 0, while ΔS (for process 2) $\neq 0$
 - (b) $q_{cyclic} = 0$ for process 1 and $q_{cyclic} \neq 0$ for process 2
 - (c) More heat can be converted to work in process 1 than in process 2
 - (d) More work can be converted to heat in process 1 than in process 2 $% \left({{{\bf{n}}_{\rm{c}}}} \right)$

<u>Attempt ALL the questions. Questions 31 – 44 (subjective questions) carry fifteen marks each.</u>

31. Identify reagent (P) and write the structure of products (Q, R, S and T) in the following series of reactions



- (i) What is the optical activity of the product?
- (ii) Draw the energy profile for the reaction
- (iii) Write the structure of the intermediate
- (iv) What is the effect of doubling the concentration of KOH on the rate of the reaction?
- (v) If aqueous KOH is replaced by alcoholic KOH, write the structure of the product formed.
- 33. (a) Suggest a method for the following transformation involving minimum number of steps.



Tiruchirappalli – 620 024

ajchemacademy@gmail.com

 \square



Indicate the reagents/reaction conditions required at each step clearly.

(b) A dipeptide on hydrolysis gives two amino acids (X) and (Y). If the dipeptide is first treated with HNO₂ and then hydrolysis is carried out, (X) and lactic acid are obtained. (X) on heating gives 2, 5-diketopiperazine as shown below. Identify (X) and (Y), and write their sequence in the dipeptide.



34. Identify the compounds A, B, C, D and E in the following reactions



35. (a) Why do the boiling points of the following compounds vary in the order,

 $H_2O > H_2Se > H_2S$

- (b) Identify the products in the reaction of CCl₄ and SiCl₄ with water. Justify your answer.
- 36. (a) Write the steps involved in the production of pure elemental silicon from silica.
 - (b) Both the products A and B, in the following reactions, contain boron and nitrogen. Identify A and B.

$$3 \text{ NH}_4 \text{CI} + 3 \text{ BCI}_3 \xrightarrow{\text{C}_6 \text{H}_5 \text{CI}} \text{A} \xrightarrow{\text{LiAIH}_4} \text{B}$$

- 37. (a) Addition of potassium oxalate solution to a hot solution of potassium dichromate containing dilute sulfuric acid leads to effervescence and formation of potassium tris-oxalatochromate(III).
 - i. Write the chemical formula of the chromium complex formed.
 - ii. Write the balanced chemical equation for the formation of the complex.
 - iii. Calculate the spin-only magnetic moment at room temperature in



JAM - 2006 - CY

Bohr magnetons of the complex.

- (b) Write the structures of possible isomers of [CoCl₂(en)₂]Cl
- **38.** (a) Draw the unit cell of CsCl lattice. Draw the (100) and (110) planes separately and indicate the positions of cesium and chloride ions.
 - (b) The hydration enthalpies of divalent metal ions of ten elements from calcium to zinc are plotted against their atomic numbers. Why do the hydration enthalpies of only three elements, Ca, Mn and Zn fall on a straight line, whereas values for other metal ions deviate from this line?



- 39. (a) 5 grams of a protein was hydrolysed into amino acids, one of which is alanine. To this mixture, 0.1 gram of partially deuterated alanine, H₂N-CH(CD₃)-COOH, was added. After thorough mixing, some of the alanine was separated and purified by crystallization. The crystalline alanine contains 0.652 weight percent of D. How many grams of alanine were originally present in 5 grams of protein?
 - (b) What is the role of ammoniacal buffer in the volumetric titration of zinc sulphate against EDTA, using Solochrome black (Eriochrome black T) indicator? Write the structure of Zn-EDTA anionic complex.
- 40. (a) Calculate the pH of a solution obtained by mixing 50.00 ml of 0.20 M weak acid HA ($K_a = 10^{-5}$) and 50.00 ml of 0.20M NaOH at room temperature.
 - (b) One mole of a salt of type MX is dissolved in 1.00 kg of water. The freezing point of the solution is -2.4 °C. Calculate the percent dissociation of the salt in water.
- 41. (a) The rate constant of the reaction $ClO + NO \rightarrow Cl + NO_2$ varies with



JAM - 2006 - CY

temperature as:

T (K)	200	400
$k (cm^{-3}s^{-1})$	$2.0 imes 10^{-11}$	$1.0 imes 10^{-11}$

Determine the Arrhenius activation energy for the reaction, assuming that the frequency factor does not change in this temperature range.

(b) Ozone seems to be formed in the atmosphere through the photolysis of diatomic molecule:

$$O_2 \xrightarrow{k_0} 2O$$

$$O_1 + O_2 \xrightarrow{k_1} O_3$$

$$O_3 + O \xrightarrow{k_2} 2O_2$$

Applying steady-state approximation, determine the rate law for the formation of ozone. Show that the formation of ozone follows first order kinetics when the concentration of O_3 is extremely small.

- 42. (a) The reaction $N_2 + 3H_2 \rightarrow 2NH_3$ is carried out at 300 K by mixing N_2 and H_2 . The standard free energy of formation of NH_3 is -16.4 kJ/mol. After one hour of mixing, the partial pressures of N_2 , H_2 and NH_3 are 50 bars, 2 bars and 200 bars, respectively. What is the reaction free energy at this stage of the reaction?
 - (b) Plot schematically the concentration dependence of molar conductivity of a strong electrolyte and a weak electrolyte in the same figure. The limiting ionic molar conductivities of K⁺ and Cl⁻ are 73.5 and 76.5 S cm²mol⁻¹, respectively. If the molar conductivity of 0.1 M KCl solution is 130.0 S cm² mol⁻¹, calculate the Kohlrausch's constant for KCl solution.
- 43. (a) The electronic wavefunction (Ψ) for hydrogen atom in the 2s state is given as

$$\Psi \propto \left[2 - \frac{r}{a_0}\right] \exp\left[-\frac{r}{2a_0}\right]$$

Determine the most probable radial distance for the electron in this state and also the position of the node (in terms of a_0).

(b) Calculate the wavelength corresponding to the lowest energy excitation of an electron confined to a one-dimensional box of length 1 nm. (Energy levels of a particle in a one-dimensional box are given by $E_n = n^2 h^2 / 8ma^2$).

Tiruchirappalli – 620 024





- 44. (a) Solve the differential equation y'' 5y' + 6y = 0 with the initial conditions y(0) = -1 and y'(0) = 0. Here y' and y'' refer to the first and the second derivatives, respectively, of y with respect to x. Verify your answer.
 - (b) For a particle with position r = 2i 3j + k and momentum p = i + 2j - 2k in m and kg ms⁻¹, respectively, calculate the magnitude of the angular momentum $L = r \times p$.

Q.No	Ans	Q.No	Ans		Q.No	Ans		Q.No	Ans
1.	С	9.	С		17.	С		25.	b
2.	С	10.	d	19	18.	d		26.	d
3.	С	11.	а	TE	19.	а		27.	b
4.	b	12.	b	GR	20.	b		28.	С
5.	d	13.	С	K	21.	а		29.	b
6.	а	14.	d		22.	b	5	30.	С
7.	а	15.	а		23.	С	SI		
8.	d	16.	b	XTC	24.	С	R		
		0		K IN	41-5				

Answer Key

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © Inspite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.
- © இந்த வினாத்தாளின் எந்தப் பகு<mark>தியும் ஏஜே கெம் அகாடமியி</mark>ன் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

