


JAM – 2008 – Chemistry



- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

www.csircoaching.com


Features

- | | |
|-------------------------------|----------------------------------|
| ➤ 300 ++ Live Classes | ➤ A Well-Defined Curriculum |
| ➤ 200 ++ Concept Wise Tests | ➤ A Strong Subject Foundation |
| ➤ 50 ++ Chapter Wise Tests | ➤ A Refined Learning Methodology |
| ➤ 50 ++ Model Tests | ➤ Updated Study materials |
| ➤ 2000 ++ Problem Discussions | ➤ Freshers Can easily understand |
| ➤ Recorded Videos | ➤ Question banks |




Attempt ALL the questions. Q.1 – Q.30 Multiple Choice Question (MCQ), carry THREE marks each (for each wrong answer: -1).


1. The correct statement describing the relationship between is



X




Y




Z


- (a) X and Y are resonance structures and Z is a tautomer.
 (b) X and Y are tautomers and Z is a resonance structure.
 (c) X, Y and Z are all resonance structures.
 (d) X, Y and Z are all tautomers.
2. Among the following, the correct statement concerning the optical activity is:
- (a) A molecule containing two or more chiral centres is always optically active.
 (b) A molecule containing just one chiral centre is always optically active
 (c) A molecule possessing alternating axis of symmetry is optically active.
 (d) An optically active molecule should have at least one chiral centre.
3. The correct order of acidity among is




(i)



(ii)




(iii)




(iv)

- (a) (i) < (ii) < (iii) < (iv) (b) (iv) < (iii) < (i) < (ii)
 (c) (ii) < (i) < (iii) < (iv) (d) (ii) < (iv) < (i) < (iii)
4. The major product obtained in the following reaction is,




- (a) (b) (c) (d)

5. The major product of the following reaction



- (a)
- (b)
- (c)
- (d)

6. The major product obtained in the following reaction is




- (a)
- (b)
- (c)
- (d)

7. **R-(–)-2-Bromooctane** on treatment with aqueous KOH mainly gives 2-octanol that is:


- (a) optically active with 'R' configuration (b) optically active with 'S' configuration
(c) a racemic mixture (d) a meso compound

8. The major product obtained in the following reaction is,




- (a)


(b)




(c) An equimolar mixture of







(d)




9. The major product obtained in the following reaction is,



10. The products of the following reaction are,






11. When one mole of ice is converted to water at 0°C and 1 atm, the work done (**L atm**) is:
 (a) 1.1×10^{-4} (b) 2.0×10^{-3} (c) 2.0×10^{-4} (d) 1.1×10^{-5}
12. When 100 g of water is reversibly heated from 50°C to 75°C at 1 atm, the change in entropy (JK^{-1}) of the universe is:
 (a) -0.31 (b) 0.31 (c) 0 (d) 3.1
13. For a zero-order reaction, **units of the rate constant** is expressed as
 (a) M^1s^{-1} (b) M^0s^{-1} (c) $\text{M}^{-1}\text{s}^{-1}$ (d) M^0s^0
14. 1×10^{-6} moles of the enzyme carbonic anhydrase dehydrates H_2CO_3 to produce 0.6 mol of CO_2 per second. **The turnover number** of the enzyme is:
 (a) $N_A \times 6 \times 10^{-5}$ (b) $(1/6) \times 10^{-5}$ (c) $(6 \times 10^5)/N_A$ (d) 6×10^5
15. Given that the most probable speed of oxygen gas is 1000 ms^{-1} , the **mean/average speed** (ms^{-1}) under the same conditions is:
 (a) 1224 (b) 1128 (c) 886 (d) 816
16. If the electrons were **spin $3/2$ particles, instead of spin $1/2$** , then the number of electrons that can be accommodated in a level are
 (a) 2 (b) 3 (c) 4 (d) 5
17. **For a particle in a cubic box**, the total number of quantum numbers needed to specify its state are
 (a) 1 (b) 2 (c) 3 (d) 9
18. The maximum number of phases that can co-exist in equilibrium for a **one component system** is:
 (a) 1 (b) 2 (c) 3 (d) 4
19. **With increasing pressure**, the temperature range over which the liquid state is stable.
 (a) Decreases (b) Increases
 (c) Remains constant (d) Decreases till the critical pressure and then increases.
20. The **conductance at infinite dilution** follows the order
 (a) $\text{Li}^+ > \text{Na}^+ > \text{K}^+$ (b) $\text{Na}^+ > \text{Li}^+ > \text{K}^+$
 (c) $\text{K}^+ > \text{Li}^+ > \text{Na}^+$ (d) $\text{K}^+ > \text{Na}^+ > \text{Li}^+$

21. The **V-shape of SO_2** is due to the presence of
 (a) two σ - and one π - bonds
 (b) two σ - and two π - bonds
 (c) two σ - bonds and one lone pair of electrons
 (d) two σ - and two π - bonds, and one lone pair of electrons
22. The correct order of the mean **bond energies** in the binary hydrides is
 (a) $\text{CH}_4 > \text{NH}_3 > \text{H}_2\text{O} > \text{HF}$
 (b) $\text{NH}_3 > \text{CH}_4 > \text{H}_2\text{O} > \text{HF}$
 (c) $\text{HF} > \text{H}_2\text{O} > \text{CH}_4 > \text{NH}_3$
 (d) $\text{HF} > \text{H}_2\text{O} > \text{NH}_3 > \text{CH}_4$
23. In **CsCl structure**, the number of **Cs^+ ions** that occupy second nearest neighbour locations of a **Cs^+ ion** is:
 (a) 6 (b) 8 (c) 10 (d) 12
24. In the given process, **X** is : ${}^{234}_{92}\text{U} \rightarrow {}^{230}_{90}\text{Th} + \text{X}$
 (a) α particle (b) β particle (c) β^+ emission (d) γ emission
25. For tetrahedral complexes, which always exhibit high spin states, the maximum **CFSE (crystal field stabilization energy)** is:
 (a) -8 Dq (b) -12 Dq (c) -16 Dq (d) -20 Dq
26. The **most abundant element** in earth's crust is:
 (a) Aluminium (b) Iron (c) Silicon (d) Oxygen
27. **Metal-carbon multiple bonds** in metal carbonyl are preferably identified from the stretching frequency of
 (a) Carbon-oxygen bond (b) Metal-carbon bond
 (c) Metal-oxygen bond (d) Carbon-carbon bond
28. In general, **magnetic moment of paramagnetic complexes** varies with temperature as
 (a) T^2 (b) T (c) T^{-2} (d) T^{-1}
29. The compound having an **S-S single bond** is:
 (a) $\text{H}_2\text{S}_2\text{O}_3$ (b) $\text{H}_2\text{S}_2\text{O}_4$ (c) $\text{H}_2\text{S}_2\text{O}_7$ (d) $\text{H}_2\text{S}_2\text{O}_8$
30. In a reaction, **$\text{Na}_2\text{S}_2\text{O}_3$ is converted to $\text{Na}_2\text{S}_4\text{O}_6$** . The equivalent weight of **$\text{Na}_2\text{S}_2\text{O}_3$** for this reaction is (**mol. wt. of $\text{Na}_2\text{S}_2\text{O}_3 = M$**)
 (a) M (b) $M/4$ (c) $M/2$ (d) $M/3$




Attempt ALL the questions. Questions 31 – 44 (subjective questions) carry fifteen marks each.


31. (a) Identify A, B and C in the following reaction sequence.




- (b) Identify D in the following reaction and suggest a suitable mechanism for its formation.



32. (a) Explain with the help of mechanisms, the observed stereo selectivity in the following epoxide formation reactions.




- (b) Explain on the basis of conformational analysis why (1R, 2S)-1, 2-dimethylcyclohexane is optically inactive at room temperature.
33. (a) Identify E, F and G in the following synthetic transformation:




- (b) An optically active compound H (C₅H₆O) on treatment with H₂ in the presence of Lindlar's catalyst gave a compound I (C₅H₈O). Upon hydrogenation with H₂ and Pd/C, compound H gave J (C₅H₁₂O). Both I and J were found to be optically inactive. Identify H, I and J.
34. (a) A disaccharide K gives a silver mirror with Tollen's reagent. Treatment of K with MeOH/HCl gives a monomethyl derivative L, which does not react with Tollen's reagent. Methylation of K with Me₂SO₄ and NaOH affords an octamethyl derivative of K, which upon acidic hydrolysis gives a 1:1 mixture of 2,3,4,6-tetra-O-methyl-D-glucose and 2,3,4-tri-O-methyl-D-glucose. Disaccharide K is also hydrolysed by the enzyme maltase. Identify K and L with proper stereochemistry.

(b) Identify M and N in the following reaction sequence.



35. In the following reaction sequence, identify P, R and S. Suggest suitable mechanism for the conversion of P → Q and R → S



36. (a) Consider the reactions.



Unstable



(i) Identify A and B.

(ii) What is the role of H_2O_2 in (I) and how does A favour the formation of Cr^{3+} ?


(iii) What is the role of H_2O_2 in (II) and how does B favour the formation of CrO_4^{2-} ?

(b) With the help of equations, illustrate the role of a cis-1, 2-diol in the titration of boric acid with sodium hydroxide.

37. (a) Draw the structure of anionic Ca(II)-EDTA chelate. How many rings are formed in the chelate and specify the number of atoms in each ring?


(b) Based on VSEPR theory draw the most stable structure of ClF_3 and XeF_4 .

38. (a) Identify A, B and C in the following reaction scheme



(b) From the Ellingham diagram given below, identify the metal oxide that can be

reduced at a lower temperature by carbon. Justify.



39. (a) For the complexes $[\text{FeF}_6]^{3-}$ and $[\text{Fe}(\text{CN})_6]^{3-}$.
- Show the hybridization using VB (valence bond) theory
 - Calculate the CFSE (crystal field stabilization energy)
- (b) Identify the dark blue complex formed when $[\text{Fe}(\text{CN})_6]^{3-}$ is treated with FeSO_4 and account for the origin of its colour.
40. (a) Consider the equilibrium,
- $$\text{A (g)} \rightleftharpoons \text{B (g)} + \text{C (g)}$$
- At a constant pressure of 1 atm, A dissociates to the extent of 50% at 500 K.
Calculate ΔG^0 (kJ mol^{-1}) for the reaction.
- (b) Consider the following redox system.
- $$\text{Q} + 2\text{H}^+ + 2\text{e}^- \rightleftharpoons \text{QH}_2 \quad E^0 = 0.699 \text{ V}$$
- Calculate the pH of the solution at 298 K, if the redox potential of the system is 0.817 V.
41. (a) A stream of oxygen molecules at 500 K exits from a pin-hole in an oven and strikes a slit that selects the molecules travelling in a specific direction. Given that the pressure outside the oven is 2.5×10^{-7} atm, estimate the maximum distance at which the slit must be placed from the pin-hole, in order to produce a collimated beam of oxygen. (Radius of $\text{O}_2 = 1.8 \times 10^{-10}$ m)
- (b) Liquid water is to be circulated to transfer heat from a source to a sink at 1 atm. Considering this arrangement as a Carnot engine, calculate the maximum theoretical efficiency that can be expected from the system.
42. (a) Using Heisenberg's uncertainty principle, derive an expression for the approximate ground state energy of a particle of mass m in a one-



dimensional box of length L.

- (b) The rate of a chemical reaction doubles when the temperature is changed from 300 K to 310 K. Calculate the activation energy (kJ mol^{-1}) for the reaction.

43. (a) Consider the reaction.



Assuming ideal behaviour, calculate ΔU^0 when 1 mol of CH_4 is completely oxidized at STP.

- (b) A photochemical reaction was carried out using monochromatic radiation (490 nm) of intensity 100 W. When the sample was irradiated for 30 min, 0.3 mol of the reactant was decomposed. Estimate the quantum efficiency assuming 50% absorption.

44. (a) Given that

$$C_p - C_v = \frac{\alpha^2 TV}{k_T}; \text{ Where } \alpha = \frac{1}{V} \left(\frac{\partial V}{\partial T} \right)_p \text{ and } k_T = -\frac{1}{V} \left(\frac{\partial V}{\partial P} \right)_T$$

For a pure substance, show that $C_p - C_v = R$ for 1 mol of an ideal gas.

- (b) Find the eigenvalues of the following 3×3 matrix given that 2 is one of the eigenvalues. Compute the determinant of the matrix using the eigenvalues.

$$\begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & -2 \\ 1 & -1 & 1 \end{bmatrix}$$

Answer Key

Q.No	Ans		Q.No	Ans		Q.No	Ans		Q.No	Ans
1.	a		9.	c		17.	c		25.	b
2.	b		10.	a		18.	c		26.	d
3.	b		11.	b		19.	b		27.	a
4.	c		12.	c		20.	d		28.	d
5.	d		13.	a		21.	d		29.	b
6.	a		14.	d		22.	d		30.	a
7.	b		15.	b		23.	b			
8.	b		16.	c		24.	a			

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © In spite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So we do not take any legal responsibility for them. If they are brought to our notice, corrections will be

done in next edition.

- © இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

