

JAM – 2010 – Chemistry



- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

www.csircoaching.com

Features

➢ 300 ++ Live Classes	A Well-Defined Curriculum
\succ 200 ++ Concept Wise Tests	A Strong Subject Foundation
\succ 50 ++ Chapter Wise Tests	> A Refined Learning Methodology
\succ 50 ++ Model Tests	Updated Study materials
\succ 2000 ++ Problem Discussions	Freshers Can easily understand
Recorded Videos	Question banks



 \bigoplus

Attempt ALL the questions. Q.1 – Q.30 Multiple Choice Question (MCQ), carry THREE marks each (for each wrong answer: –1).

- 1. The molar internal energy of a gas at temperature T is $U_m(T)$. The molar internal energy at T = 0 is $U_m(0)$. The correct expression that relates these two with appropriate contributions is:
 - (a) $U_m(T) = U_m(0) + 3RT$ [linear molecule; translation only]
 - (b) $U_m(T) = U_m(0) + \frac{5}{2}RT$ [linear molecule; translation and rotation only]
 - (c) $U_m(T) = U_m(0) + \frac{3}{2}RT$ [non-linear molecule; translation and rotation only]
 - (d) $U_m(T) = U_m(0) + RT$ [non-linear molecule; translation only]
- 2. If a particle has linear momentum $\vec{p} = -2\vec{i} + \vec{j} + \vec{k}$ at position $\vec{r} = 3\vec{i} \vec{j} + \vec{k}$, then its angular momentum is:

(a)
$$\vec{i} + 2\vec{k}$$
 (b) $-2\vec{i} - 5\vec{j} + \vec{k}$ (c) $5\vec{i} - 2\vec{j}$ (d) $2\vec{i} + 5\vec{j} - \vec{k}$

- 3. If Ψ is the eigenfunction to the Hamiltonian operator with α as the eigenvalue, then α MUST be
 - (a) Positive (b) Negative (c) An integer (d) Real
- 4. A quantum mechanical particle of mass m free to rotate on the surface of a sphere of radius r is in the state with energy $\frac{10\hbar^2}{mr^2}$. The degeneracy of this state is: (a) 20 (b) 10 (c) 9 (d) 4
- 5. Choose the **INCORRECT** statement among the following:
 - (a) When ideal gases are mixed, the entropy of mixing is always positive.
 - (b) At equilibrium, the chemical potential of a species is the same in all of the phases of the system.
 - (c) The total pressure of a mixture of ideal gases is equal to the sum of the partial pressure of each gas in the mixture.
 - (d) When a gas is allowed to expand, the maximum work is obtained when the process is carried out irreversibly.
- 6. The work done during the free expansion of one mole of an ideal gas at 27 °C to twice its original volume is (given: $RT = 2494 \text{ J mol}^{-1}$, $\ln 2 = 0.7$, $\log 2 = 0.3$)

iol ⁻¹
1

7. Choose the correct order of the diffusion coefficients of the following at 298 K.

H ⁺ in water	OH⁻ in water	H ₂ O in water	Sucrose in water
Р	Q	R	S



(a)	Р	>	Q	>	R	>	S	(t	b)	S	>	R	>	Q	>	Р

- (c) S > Q > R > P (d) P > R > Q > S
- 8. Two matrices are given as $X = \begin{pmatrix} 1 & 5 \\ 3 & 7 \end{pmatrix}$ and $Y = \begin{pmatrix} 2 & 4 \\ 6 & 0 \end{pmatrix}$. If X^T is the transpose of X then what would be X^TY ?
 - (a) $\begin{pmatrix} 20 & 52 \\ 4 & 20 \end{pmatrix}$ (b) $\begin{pmatrix} 20 & 4 \\ 52 & 20 \end{pmatrix}$ (c) $\begin{pmatrix} 32 & 4 \\ 48 & 12 \end{pmatrix}$ (d) $\begin{pmatrix} 44 & 28 \\ 12 & 12 \end{pmatrix}$
- 9. Addition of 1.0 g of a compound to 10 g of water increases the boiling point by 0.3 °C. The amount of compound needed to prepare a 500 ml of 0.1 M solution is (given: assume negligible dissociation or association of the compound, boiling point constant K_b of water = 0.513 K kg mol⁻¹)
 - (a) 0.855 g (b) 17.1 g (c) 8.55 g (d) 85.5 g
- 10. The molar conductivity of 0.009 M aqueous solution of a weak acid (HA) is 0.005 S m² mol⁻¹ and the limiting molar conductivity of HA is 0.05 S m² mol⁻¹ at 298 K. Assuming activity coefficients to be unity, the acid dissociation constant (K_a) of HA at this temperature is:
 (a) 1 × 10⁻⁴
 (b) 0.1
 (c) 9 × 10⁻⁴
 (d) 1.1 × 10⁻⁵
- **11.** The colour of potassium dichromate is due to 930¹
 - (a) d-d transition (b) Transition in K⁺ ion.
 - (c) Ligand to metal charge transfer (d) Metal-to-ligand charge transfer.
- 12. Which one of the following configuration will show Jahn-Teller distortion in an octahedral field?

(a) High spin d^8 (b) High spin d^4 (c) High spin d^5 (d) Low spin d^6 .

13. B_2H_6 and B_4H_{10} , respectively, are examples of

- (a) Nido and arachno boranes (b) Nido and closo boranes
- (c) Closo and arachno boranes (d) Nido boranes.
- 14. Which of the following has a square planar shape according to the VSEPR theory?

		(Atomic number:	B = 5, S = 16, Xe = 54)
(a) XeO_2F_2	(b) SF ₄	(c) BF ₄	(d) XeF ₄

15. The structure of rock salt consists of

(a) A cubic close-packed array of anions with cations in all the octahedral sites.

- (b) A cubic close-packed array of cations with anions in all the tetrahedral sites.
- (c) A hexagonal close-packed array of anions with cations in all the octahedral sites.



JAM - 2010 - CY



(d) A cubic close-packed array of anions with cations in all the tetrahedral sites Among lithium, nitrogen, carbon and oxygen, which element has the highest first 16. ionization potential? (a) Lithium (b) Nitrogen (c) Carbon (d) Oxygen In which of the following C–H bond has the highest 's' character? 17. (a) Acetylene (b) Ethylene (c) Methane (d) CH radical 18. Which one of the following is an electron deficient molecule according to the octet rule? (a) CH₄ (b) $H_3N: BH_3$ (c) AlH_3 (d) GeH₄ 19. Which one of the following has the highest lattice energy? CATE(c) LiF (b) $CaCl_2$ (a) LiCl (d) KCl 20. At room temperature, HCl is a gas while HF is a liquid because (a) of a strong bond between H and F in HF (b) HF is less acidic as compared to HCl (c) of strong intermolecular H-bonding in HF (d) HCl is less acidic as compared to HF **Benzene and Dewar benzene are** 21. Benzene **Dewar benzene** (b) Structural isomers (a) Canonical forms (d) Conformational isomers (c) Tautomers 22. The **IUPAC** name of the following compound is: CI CN CH-CH H₃C CH₃ (a) 2-cyano-3-chlorobutane (b) 2-chloro-3-cyanobutane (d) 3-chloro-2-methylbutanenitrile (c) 2-methyl-3-chlorobutanenitrile 23. Which chemical test will distinguish the compounds shown below? $-CH_3$ -CH₂Br Br (b) Ethanolic silver nitrate test (a) Beilstein's flame test (c) Sodium fusion test (d) Fehling's test

0

ajchemacademy@gmail.com \square



JAM-2010-CY

24. The reaction of the bromo compound shown below with sodium ethoxide gives predominantly



The position in the graph that corresponds to the isoelectric point of alanine is:

27. The absolute configurations at the two chiral centers in **D-Ribulose** are



JAM - 2010 - CY



28. The most stable conformation of the molecule shown below is correctly represented by



29. Thermal rearrangement of the following compound would give



30. The energy profile diagram that corresponds to **1**, **2**-dihydroxyethane for rotation around the C–C bond is







<u>Attempt ALL the questions. Questions 31 – 44 (subjective questions)</u> <u>carry fifteen marks each.</u>

- 31. (a) Equilibrium constant for a reaction doubles as the temperature is increased from 300 K to 600 K. Calculate the standard reaction enthalpy (in kJ mol⁻¹) assuming it to be constant in this temperature range.
 - (given R = 8.3 J K⁻¹ mol⁻¹, ln 2 = 0.7).
 (b) A 50 mL solution of 0.1 M monoprotic acid (K_a = 1 × 10⁻⁵ at 298 K) is titrated with 0.1 M NaOH at 298 K. Calculate the [H⁺] of the solution after the addition of 50 mL of NaOH at this temperature. (given K_w = 1 × 10⁻¹⁴ at 298 K)
- 32. For the reaction, $H_2(g) + Br_2(g) \rightarrow 2HBr(g)$ the following mechanism has been proposed.

CADEM

Initiation: (M is the initiator/terminator)

 $Br_2 + M \xrightarrow{k_i} Br + Br + M$ **Propagation:**

$$\begin{array}{c} \stackrel{\odot}{\text{Br}} + \text{H}_2 \xrightarrow{k_p} \text{HBr} + \stackrel{\odot}{\text{H}} \\ \stackrel{\odot}{\text{H}} + \text{Br}_2 \xrightarrow{k_{p'}} \text{HBr} + \stackrel{\odot}{\text{Br}} \end{array}$$

Retardation:

$$H^{\odot}$$
 + HBr $\xrightarrow{k_r}$ H_2 + Br

Termination:

 $B_r^{\odot} + B_r^{\odot} + M \xrightarrow{k_t} B_r^{\circ} + M + energy$

- (a) Write the differential rate equations for the formation of the two intermediates $H^{\odot}_{and} Br^{\odot}_{and}$.
- (b) Using the steady-state approximation, calculate the concentration of the



intermediates $\overset{\frown}{H}^{\odot}$ and $\overset{\Box}{Br}^{\odot}$ and obtain the rate law for the formation of HBr. Calculate ΔH_m and ΔS_m for the process



Assume that at 273 K the molar enthalpy of fusion of ice is 6006 J mol^{-1} , the heat capacity $C_{p,m}(s)$ of ice is 38 J $K^{-1}mol^{-1}$ and heat capacity $C_{p,m}(l)$ of liquid water is 76 J K^{-1} mol⁻¹. Consider the heat capacities to be constants.

(given: ln 263 = 5.57 and ln 273 = 5.61)

7

Two beakers, one containing 0.02 M KMnO₄, 0.2 M MnSO₄ and 0.5 M H₂SO₄ and **34**. another containing 0.15 M FeSO₄ and 0.05 M Fe₂(SO₄)₃, are connected by a saltbridge. Platinum electrodes are placed in each beaker and these two electrodes are connected via a wire with a voltmeter in between. H_2SO_4 is present in equal volumes in each beaker. Assume H₂SO₄ is completely ionized.

Given: $E_{Fe^{3+}/Fe^{2+}}^{0} = 0.8 V$, $E_{MnO_{4}^{-}/Mn^{2+}}^{0} = 1.5 V$, $\frac{2.303 RT}{F} = 0.06V$ and $\log 2 = 0.3$

- (a) Write the complete balanced redox reaction for this cell.
- (b) What would be the potential of each half-cell after the reaction has reached equilibrium?
- An atomic orbital is described by the wavefunction 35. GENE ACA

$$\phi(\mathbf{r}) = \frac{1}{\sqrt{\pi a_0^3}} e^{-\left(\frac{\mathbf{r}}{a_0}\right)}$$
, where a_0 is the Bohr radius.

Given: $d\tau = r^2 \sin\theta \, dr \, d\theta \, d\phi$ and $\int_0^\infty r^n e^{-\beta r} dr = \frac{n!}{\beta^{n+1}}$ (n is a positive integer)

- (a) Identify the atomic orbital and calculate the mean or the average radius of this orbital in terms of a_0 .
- (b) Calculate the most probable radius (in terms of a_0) at which an electron Will be found when it occupies this orbital.
- **36**. Identify W, X, Y and Z in the following sequence

$$Li + W(g) \xrightarrow{heat} X \xrightarrow{H_20} Y(g) \xrightarrow{alkaline K_2HgI_4} Z$$

$$(X = red) \qquad (Z = brown)$$

Y turns moist litmus paper blue. Write balanced chemical equation for the



33.

conversion of Y to Z.

37. (a) Draw the crystal field splitting diagram with appropriate labels for [NiCl₄]²⁻.
 Determine the spin only magnetic moment and the crystal field stabilization energy (CFSE) for this complex.

(given: atomic number of Ni = 28)

8

- (b) Write the balanced equations for the reactions involved in the iodometric estimation of Cu^{2+} using thiosulfate.
- **38.** (a) In the reaction sequence given below P is an anionic Fe(II) complex.

$$P \xrightarrow{aq.NO_2^-} Q \xrightarrow{aq.S^{2-}} R$$

$$(Q = brown) \qquad (R = purple)$$

Identify P, Q and R. 🐰 💆

- (b) Draw a properly labeled unit cell diagram of CsCl. Show through calculations that there is only one CsCl per unit cell.
- 39. (a) Write the balanced chemical equations for the reactions involved in the synthesis of borazine using ammonium chloride as one of the starting materials. Write the structure of borazine.
 - (b) Draw Lewis structures of SF₄ and NO₃⁻
- 40. (a) Complete the following sequence by identifying **E**, **F** and **G**.



(b) Identify H and I in the reactions below.



41. (a) Identify the products J, K, and L in the following reactions. Lassigne's test for L shows the presence of nitrogen only.



(b) Write the structure of **M** and **N** in the following reactions.

ajchemacademy@gmail.com



JAM-2010-CY

$$\begin{array}{c|c} & Me \\ CH_2C \equiv CH \\ \hline \\ O \\ \end{array} \begin{array}{c} HgSO_4 \\ \hline \\ dilute H_2SO_4 \end{array} M \begin{array}{c} NaOEt/EtOH \\ \hline \\ heat \end{array} N$$

42. (a) Write the structures of **P**, **Q** and **R** in the given reaction sequence.

$$Ph \longrightarrow H \xrightarrow{1. MeMgBr} P \xrightarrow{Pd/BaSO_4} Q \xrightarrow{LiAlH_4} R$$

(b) Identify **S** and **T** in the reactions given below:

43. (a) Identify X, Y and Z in the following reactions



(b) Suggest a suitable mechanism for the following reaction.



44. Consider the following reactions for a compound with molecular formula $C_{10}H_{16}$.



- (a) Write structures that are consistent with the above data for the formula $C_{10}H_{16}$.
- (b) Given that myrcene is a terpene and has the molecular formula $C_{10}H_{16}$, using the isoprene rule identify the correct structure for myrcene among the structures elucidated in part (a).

Tiruchirappalli – 620 024



Q.No	Ans	Q.No	Ans	Q.No	Ans	Q.No	Ans
1.	b	9.	С	17.	а	25.	а
2.	b	10.	а	18.	С	26.	b
3.	d	11.	С	19.	b	27.	а
4.	С	12.	b	20.	С	28.	d
5.	d	13.	а	21.	b	29.	С
6.	С	14.	d	22.	d	30.	С
7.	а	15.	а	23.	b		
8.	b	16.	b	24.	С		

Answer Key

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- Inspite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.

© இந்த வினாத்தாளின் எ<mark>ந்தப் பகுதியும் ஏஜே கெம் அகாடமியின்</mark> முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.

© இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

