

JAM – 2011 – Chemistry



www.csircoaching.com

- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

Features							
۶	300 ++ Live Classes	۶	A Well-Defined Curriculum				
≻	200 ++ Concept Wise Tests	۶	A Strong Subject Foundation				
\triangleright	50 ++ Chapter Wise Tests	۶	A Refined Learning Methodology				
\triangleright	50 ++ Model Tests	≻	Updated Study materials				
≻	2000 ++ Problem Discussions	\triangleright	Freshers Can easily understand				
≻	Recorded Videos	≻	Question banks				



Attempt ALL the questions. Q.1 – Q.30 Multiple Choice Question (MCQ), carry THREE marks each (for each wrong answer: -1). The pair of semimetals in the following is: 1. (a) Al, Si (b) Ga, As (c) Sb, Te (d) Ca, B 2. The most probable oxidation states for both Cr and Mo are (a) + 2, + 3, + 4(b) + 2, + 3, + 5(c) + 2, + 3, + 6(d) +3, +4, +53. The correct order of acidic character is: (a) $Al_2O_3 > MgO$ $> SiO_2 > P_4O_{10}$ (b) $P_4O_{10} > Al_2O_3 > MgO > SiO_2$ (c) $P_4O_{10} > SiO_2 > Al_2O_3 > MgO$ (d) $SiO_2 > P_4O_{10} > Al_2O_3 > MgO$ 4. The pair of amphoteric oxides is: (b) $V_2 O_3$, $Cr_2 O_3$ (c) VO_2 , $Cr_2 O_3$ (a) VO, Cr_2O_3 (d) $V_2 O_5$, $Cr O_3$ In the structure of $B_4O_5(OH)_4^{2-}$ 5. (a) All four B atoms are trigonal planar (b) One B atom is tetrahedral and the other three are trigonal planar. (c) Three B atoms are tetrahedral and one is trigonal planar. (d) Two B atoms are tetrahedral and the other two are trigonal planar. The **pH** of an aqueous solution of Al³⁺ is likely to be 6. (b) Acidic (c) Slightly basic (a) Neutral (d) Highly basic. Hydrolysis of (CH₃)₂SiCl₂ and CH₃SiCl₃ leads to 7. (a) Linear chain and cross-linked silicones, respectively. (b) Cross-linked and linear chain silicones, respectively. (c) Linear chain silicones only (d) Cross-linked silicones only. 8. The oxide that has the inverse spinel structure is: (a) $FeCr_2O_4$ (b) $MnCr_2O_4$ (c) $CoAl_2O_4$ (d) Fe_2CoO_4 9. The transition metal monoxide that shows metallic conductivity is: (a) NiO (b) MnO (c) TiO (d) CoO 10. The metal that is extracted by the reduction method is: (a) Al (b) Au (c) Hg (d) Mg11. The most viscous liquid is: (a) Water (b) Methanol (c) Ethylene glycol (d) Glycerol Tiruchirappalli - 620 024 www.csircoaching.com ajchemacademy@gmail.com

0

- 12. In ammoniacal buffer, oxine (8-hydroxyquinoline) forms yellow precipitate with (a) Mg^{2+} (b) Ca^{2+} (c) Ba^{2+} (d) Sr^{2+}
- 13. Addition of an aqueous solution of Fe(II) to potassium hexacyanochromate (III) produces a brick-red colored complex, which turns dark green at 100 °C. The dark green complex is:
 - (a) $Fe_4[Cr(CN)_6]_3$ (b) $KFe[Cr(CN)_6]$ (c) $KCr[Fe(CN)_6]$ (d) $Fe[Cr(CN)_6]$
- 14. In the following equation X is:

(a)
$$2 \frac{1}{95}Am + \alpha \rightarrow \frac{243}{97}Bk + X$$

(b) $\frac{1}{95}n$ (c) $2 \frac{1}{1}H$ (d) $\frac{4}{2}He$

- 15. Based on the principle of equipartition of energy, the molar heat capacity of CO_2 at GATE & JAN constant volume C_{v.m} is:
 - (b) 6R (d) 9R (a) 3.5 R (c) 6.5R
- 16. One mole of a van der Waals gas undergoes reversible isothermal transformation from an initial volume V_1 to a final volume V_2 . The expression for the work done is:

(a) RT
$$\ln \frac{V_2}{V_1} + a(V_2 - V_1)$$

(b) $-RT \ln \frac{V_2 - b}{V_1 - b} + a(\frac{1}{V_1} - \frac{1}{V_2})$
(c) RT $\ln \frac{P_2}{P_1}$
(d) RT $\ln \frac{V_2 - b}{V_1 - b} - a(\frac{1}{V_1} - \frac{1}{V_2})$

- 17. The scalar product of two vectors u and v, where u = 2i + 3j 5k and v = i + j + j3k, is
 - (b) 2i + 3j 15k (c) 3i + 4j 2k(a) – 10 (d) 10
- 18. The minimum concentration of silver ions that is required to start the precipitation of $Ag_2S(K_{sp} = 1 \times 10^{-51})$ in a 0.1 M solution of S^{2-} is: (b) 1×10^{-50} M (a) 1×10^{-49} M (c) 1×10^{-26} M (d) 1×10^{-25} M
- 19. Identify the correct statement regarding Einstein's photoelectric effect
 - (a) The number of electrons ejected depends on the wavelength of incident radiation
 - (b) Electron ejection can occur at any wavelength of incident radiation.
 - (c) The number of electrons ejected at a given incident wavelength depends on the intensity of the radiation.
 - (d) The kinetic energy of the ejected electrons is independent of the wavelength of incident radiation.
- The hydrolysis constant (K_h) of NH₄Cl is 5.6 × 10⁻¹⁰. The concentration of H₃O⁺ 20. in a 0.1 M solution of NH₄Cl at equilibrium is:





(a) $\sqrt{5.6 \times 10^{-11}}$ (b) $\sqrt{5.6 \times 10^{-10}}$ (c) 5.6×10^{-10} (d) 2.8×10^{-5}

- 21. The acid dissociation constant (K_a) for HCOOH, CH₃COOH, CH₂ClCOOH and HCN at 25 °C are 1.8×10^{-4} , 1.8×10^{-5} , 1.4×10^{-3} and 4.8×10^{-10} respectively. The acid that gives highest pH at the equivalence point when 0.2 M solution of each acid is titrated with a 0.2 M solution of sodium hydroxide is:
 - (a) HCOOH (b) CH_3COOH (c) $CH_2CICOOH$ (d) HCN
- 22. For an ideal gas undergoing reversible Carnot Cycle, the plot of enthalpy (H) versus entropy (S) is:



23. Hybridizations of the atoms indicated with the asterisk (*) in the following compounds sequentially are





24. The Cahn-Ingold-Prelog (CIP) priorities of the groups and the absolute configuration (R/S) of the following compound are





(a)	CH ₂ OH	>	$CH(CH_3)_2$	>	CH=CH ₂	>	CH_3	and	S
(b)	CH ₂ OH	>	CH=CH ₂	>	$CH(CH_3)_2$	>	CH_3	and	S
(c)	CH ₂ OH	>	CH=CH ₂	>	$CH(CH_3)_2$	>	CH_3	and	R
(d)	CH ₂ OH	>	$CH(CH_3)_2$	>	CH=CH ₂	>	CH₃	and	R

25. The optically active stereoisomer of the following compound is:







	(i)		(ii)		(iii)		(iv)
(a)	Identical	,	enantiomers	,	diastereomers	and	structural isomers
(b)	enantiomers	,	Identical	,	structural isomers	and	diastereomers
(c)	enantiomers	,	Identical	,	diastereomers	and	structural isomers
(d)	Identical	,	Identical	,	diastereomers	and	structural isomers

28. The **INCORRECT** statement in the following is:

- (a) The nucleobase pairs are aligned perpendicular to the helical axis in DNA.
- (b) RNA contains uracil and thymine, but DNA contains only thymine.
- (c) All naturally occuring amino acids with the exception of glycine are chiral
- (d) All enzymes are proteins, but all proteins are not necessarily enzymes.

29. The products P and Q in the following reactions, respectively, are



30. The major product in the following reaction is:



<u>Attempt ALL the questions. Questions 31 – 44 (subjective questions)</u> <u>carry fifteen marks each.</u>

31. (a) In the following reactions, identify X, Y and Z.

 $Na_2SO_3 + S \xrightarrow{\text{boiling water}} X$ (colourless solid)

Tiruchirappalli – 620 024





JAM-2011-CY

AgBr
$$\xrightarrow{\text{excess X}}$$
 Y (soluble complex)
X + Cl₂ + H₂O $\xrightarrow{\text{boiling water}}$ Z + HCl

(b) Draw the structures of $S_4N_4H_4$ and $N_4S_4F_4$.

- 32. (a) The magnetic moment of [Fe(Phen)₂(NCS)₂] varies with temperature. The magnetic moments at 200 K and 50 K are 4.9 BM and 0 BM, respectively. Write the d-electron configurations of Fe at both temperatures and give reason for the observed change in the magnetic moment. (phen = 1,10-phenanthroline)
 - (b) PCl₅ exists as a discrete covalent molecule in the gaseous state, but is ionic in the solid state. Draw the structures of PCl₅ in gaseous and solid states.
- 33. In the following equilibrium and reactions, identify species **B** to **E**. Write the balanced chemical equation for the conversion of **C** to **E**.



- $B + diphenylcarbazide \rightarrow D$ (violet color)
- $C + HCl \rightarrow E$ (greenish yellow gas)
- 34. (a) Identify species A and C in the following.

Write the balanced chemical equation for the conversion of A to A^{3+} .

 \mathbf{A} + aquaregia $\rightarrow \mathbf{A}^{3+}$ + NO

 $A^{3+} + I^- \rightarrow B$ (black precipitate)

 $\mathbf{B} + \mathbf{I}^{-}(\text{excess}) \rightleftharpoons \mathbf{C}$ (orange color)

Hint: C on the dilution with water gives B

- (b) Draw the structures of X and Y in the following reactions.
 - (i) Borazine + HCl \rightarrow X
 - (ii) Borazine + $Br_2 \rightarrow Y$
- 35. (a) The molar conductances at infinite dilution for BaCl₂, KCl, K₂SO₄ and Cl⁻ are 280, 150, 300 and 76 Ω^{-1} m² mol⁻¹, respectively. Calculate the transport number of Ba²⁺ in BaSO₄ solution at infinite dilution.
 - (b) If 4 moles of a MX₂ salt in 1 kg of water raises the boiling point of water by
 3. 2 K, Calculate the degree of dissociation of MX₂ in the solution.

[For water, $K_b = 0.5 \text{ K kg mol}^{-1}$]





36. (a) For the reaction R → P, the plot of ln[R] versus time (t) gives a straight line with a negative slope. The half-life for the reaction is 3 minutes.

 $(\ln 2 = 0.693, \ln 0.1 = -2.303)$

- (i) Derive the expression for $t_{1/2}$.
- (ii) Calculate the slope of the straight line
- (iii) Calculate the time required for the concentration of R to decrease to 10% of its initial value.
- (b) Shown below is the Jablonski diagram that describes various photophysical processes. The solid arrows (→) represent radiative transitions and the wavy arrow (→) represents a non-radiative transition



- (i) Name the photophysical pathways X, Y and Z.
- (ii) Which of the radiative decays is faster?
- 37. (a) (i) Given that $\Delta G = -nFE$, derive the expression for the temperature

dependence of the cell potential (E) in terms of the change in entropy (ΔS).

(ii) For a cell reaction, E (at 25 °C) = 1.26 V, n = 2 and $\Delta S = -96.5 \text{ J K}^{-1} \text{ mol}^{-1}$. Calculate E at 85 °C by assuming ΔS to be independent of temperature.

 $(F = 96500 \text{ C mol}^{-1}).$

7

(b) The phase diagram for the lead-antimony system at a certain pressure is given below.



(i) Identify the phases and components in region-I and region-II.

(ii) Calculate the number of degrees of freedom (Variance) at point M



- 38. (a) One mole of an ideal gas initially at 300 K and at a pressure of 10 atm undergoes adiabatic expansion.
 - i. Reversibly and
 - ii. Irreversibly against a constant external pressure of 2 atm until the final pressure becomes equal to the external pressure. Calculate ΔS_{system} for (i) and (ii). For (ii), express the final answer in terms of R.

Given: Molar heat capacity at constant volume $C_{v,m} = 3R/2$

8

(b) For the following equilibrium at 300 °C.

$$N_2O_4$$
 (g) $- 2NO_2$ (g)

Calculate K_p when N_2O_4 is 30% dissociated and the total pressure is 2 bar.

39. (a) The Maxwell probability distribution of molecular speeds for a gas is:

$$\mathbf{F}(\mathbf{v})\mathbf{d}\mathbf{v} = 4\pi v^2 \left(\frac{\mathbf{m}}{2\pi \mathbf{k}T}\right)^{3/2} \exp\left(-\frac{\mathbf{m}v^2}{2\mathbf{k}T}\right) \mathbf{d}\mathbf{v}$$

Where 'v' is the speed, 'm' the mass of a gas molecule and k the Boltzmann constant.

- i. Use F(v) to show that the most probable speed v_{mp} is given by the expression. $v_{mp} = \left(\frac{2RT}{M}\right)^{\frac{1}{2}}$
- ii. Use $R = 8 \text{ J } \text{K}^{-1} \text{ mol}^{-1}$ in the above expression to calculate the v_{mp} for $CH_4(g)$ at 127 °C
- (b) The wavefunction of a quantum state of hydrogen atom with principal quantum number n = 2 is:

$$\varphi_{2\mathrm{lm}}(\mathbf{r},\boldsymbol{\theta},\boldsymbol{\phi}) = \frac{1}{\sqrt{32\pi}} \left(\frac{1}{a_0}\right)^{3/2} \left(2 - \frac{\mathbf{r}}{a_0}\right) \exp\left(-\frac{\mathbf{r}}{2a_0}\right)$$

- i. Identify the values of quantum numbers ℓ and m and hence the atomic orbital.
- ii. Find where the radial node of the wavefunction occurs.
- 40. (a) Write the possible substitution products in the following reactions. Indicate the type(s) of mechanisms $(S_N 1/S_N 2/S_N 2')$ that is/are operative in each reaction.

(i)
$$Br \xrightarrow{CN, DMF} ?$$



(ii)
$$\swarrow$$
 Br $\xrightarrow{CH_3OH}$?

(b) Write the elimination products A to C in the following reaction. Identify the **major Product**



41. (a) Write the structures of A to C in the following reaction sequence.



43. Write the structures of products A to E in the following reaction sequence.

0



(stable product)



44. Oxanamide O, a tranquilizer, is synthesized according to the following reaction scheme. Write the missing structures and reagents K to O.



Q.No Q.No Ans Q.No Ans Ans Q.No Ans 1. 9. 17. 25. b С С а 2. 10. 18. 26. С С d а 3. 11. d 19. 27. С С С 4. 12. 20. 28. b С а а 5. d 13. 21. d 29. С С b 14. 22. 6. b 30. а С 7. 15. 23. а С а 8. d 16. b 24. b

Answer Key

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © Inspite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So, we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.





- © இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.



Tiruchirappalli – 620 024

O



