


## JAM – 2014 – Chemistry



[www.csircoaching.com](http://www.csircoaching.com)

- ✓ CSIR-NET & SLET | SET Chemistry Coaching
- ✓ University Chemistry Entrance (PhD | PG)
- ✓ GATE Chemistry Coaching
- ✓ CUET-PG & JAM Chemistry Coaching

### Features

- |                               |                                  |
|-------------------------------|----------------------------------|
| ➤ 300 ++ Live Classes         | ➤ A Well-Defined Curriculum      |
| ➤ 200 ++ Concept Wise Tests   | ➤ A Strong Subject Foundation    |
| ➤ 50 ++ Chapter Wise Tests    | ➤ A Refined Learning Methodology |
| ➤ 50 ++ Model Tests           | ➤ Updated Study materials        |
| ➤ 2000 ++ Problem Discussions | ➤ Freshers Can easily understand |
| ➤ Recorded Videos             | ➤ Question banks                 |



**Attempt ALL the questions. Q.1 – Q.35 Multiple Choice Question (MCQ), carry TWO marks each (for each wrong answer: -0.66).**


- For square matrices  $M$  and  $N$ , if  $MN = M$  and  $NM = N$ , then:
  - $M^2 = M$  and  $N^2 = N$
  - $N^2 \neq N$  and  $M^2 = M$
  - $M^2 \neq M$  and  $N^2 \neq N$
  - $M^2 \neq M$  and  $N^2 = N$
- The energy of an electron in a hydrogenic atom with nuclear charge  $Z$  varies as:
  - $Z$
  - $Z^2$
  - $1/Z$
  - $1/Z^2$
- The carbonyl stretching frequency ( $\nu_{C=O}$ ) is highest for:
  - 
  - 
  - 
  -
- The homolytic breaking of the  $C_a-C_b$  bond is easiest in:
  - 
  - 
  - 
  -
- Tollen's test will be negative for:
  - Glucose
  - Mannose
  - Sucrose
  - Galactose
- Which one among the following is a sesquiterpene?
  - 
  - 
  - 
  -
- The predicted geometry of  $TeF_4$  by VSEPR theory is:
  - Octahedral
  - Square planar
  - Tetrahedral
  - Trigonal bipyramidal
- Among the following, the isoelectronic pair is:
  - $NO$  and  $CO$
  - $O_2^-$  (superoxide anion) and  $NO^-$
  - $NO^+$  and  $CO$
  - $O_2^-$  (superoxide anion) and  $NO^+$
- The metal ion of an enzyme involved in hydration of  $CO_2$  is:
  - $Cu(II)$
  - $Fe(II)$
  - $Mg(II)$
  - $Zn(II)$

10. Among the following, the element having **maximum inert pair effect** is:

[Given: Atomic number of Ge = 30, Pb = 82, Si = 14 and Sn = 50]

- (a) Ge (b) Pb (c) Si (d) Sn

11. The reactivity of compounds **I-IV** with maleic anhydride (**V**) follows the order:




- (a) I < II < III < IV (b) II < IV < III < I  
(c) II < I < III < IV (d) II < I < IV < III


12. Which one among the following molecules is **chiral**?




13. Identify the **starting material "X"** in the given reaction.




14. The major product for the following reaction is:







15. The structure of the major product in the following reaction is:



16. The **correct** orientation of dipoles in pyrrole and pyridine is:




17. Specific rotations of freshly prepared aqueous solutions of **I** and **II** are **+112** and **+18.7**, respectively. On standing the optical rotation of aqueous solution of “I” slowly decreases to give a final value of **+52.7** due to equilibration with **II**. Under this state of equilibrium, what is the ratio **II : I**?



- (a) 0.57 (b) 1.00 (c) 1.75 (d) 5.9

18. The major product formed in the following reaction is:



19. In **boron neutron capture therapy**, the initial boron isotope used and the particle generated after neutron capture respectively are:

- (a)  $^{11}\text{B}$  and  $\alpha$  particle (b)  $^{10}\text{B}$  and  $\alpha$  particle  
(c)  $^{11}\text{B}$  and  $\beta$  particle (d)  $^{10}\text{B}$  and  $\beta$  particle

20. The number of  $\alpha$  and  $\beta$  particle(s), generated in the following radioactive decay process, are:  $^{238}_{92}\text{U} \rightarrow ^{234}_{92}\text{U}$

- (a) one  $\alpha$  and two  $\beta$  particles (b) two  $\alpha$  and one  $\beta$  particles  
(c) one  $\alpha$  and four  $\beta$  particles (d) no  $\alpha$  and four  $\beta$  particles

21. In the measurement of hardness of water by complexometric titration, identify **P** and **Q** in the following equation, (**In** is Indicator)



- (a)  $\text{P} = \text{MgY}$  ;  $\text{Q} = \text{MgIn}$  (b)  $\text{P} = \text{MgY}_2$  ;  $\text{Q} = \text{MgIn}_2$   
(c)  $\text{P} = \text{MgIn}_2$  ;  $\text{Q} = \text{MgY}_2$  (d)  $\text{P} = \text{MgIn}$  ;  $\text{Q} = \text{MgY}$

22. An aqueous solution of haemoglobin has a molar absorptivity value of **18,600  $\text{L mol}^{-1} \text{cm}^{-1}$**  for an absorbance value of 0.1 at 540 nm. The concentration (in  $\mu\text{M}$ ) of the haemoglobin solution is: (Given: cell thickness = 1cm).

- (a) 0.537 (b) 5.37 (c) 53.7 (d) 537.0

23. The electronic transitions responsible for the colour of  **$\text{K}_2\text{Cr}_2\text{O}_7$**  and porphine in their solid state respectively are:



- (a)  $d \rightarrow d$  ;  $\pi \rightarrow \pi^*$  (b)  $M \rightarrow L$  charge transfer;  $\pi \rightarrow \pi^*$   
(c)  $L \rightarrow M$  charge transfer;  $\pi \rightarrow \pi^*$  (d)  $L \rightarrow M$  charge transfer;  $d \rightarrow d$

24. The correct order of **M–C** (**M = Ti, V, Cr and Mn**) bond stretching frequency is:

(Given: Atomic number of **Ti = 22, V = 23, Cr = 24 and Mn = 25**)

- (a)  $[\text{V}(\text{CO})_6]^- < [\text{Cr}(\text{CO})_6] < [\text{Mn}(\text{CO})_6]^+ < [\text{Ti}(\text{CO})_6]^{2-}$   
(b)  $[\text{Ti}(\text{CO})_6]^{2-} < [\text{V}(\text{CO})_6]^- < [\text{Cr}(\text{CO})_6] < [\text{Mn}(\text{CO})_6]^+$   
(c)  $[\text{Mn}(\text{CO})_6]^+ < [\text{Cr}(\text{CO})_6] < [\text{V}(\text{CO})_6]^- < [\text{Ti}(\text{CO})_6]^{2-}$   
(d)  $[\text{Mn}(\text{CO})_6]^+ < [\text{V}(\text{CO})_6]^- < [\text{Cr}(\text{CO})_6] < [\text{Ti}(\text{CO})_6]^{2-}$




25. For the following reactions, the metal complexes **X** and **Y** are:



- (a)  $X = \text{Ni}(\text{CO})_4$ ;  $Y = \text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2$  (b)  $X = \text{Ni}(\text{CO})_4$ ;  $Y = \text{Fe}(\eta^1\text{-C}_5\text{H}_5)_2$   
 (c)  $X = \text{Ni}(\text{CO})_5$ ;  $Y = \text{Fe}(\eta^5\text{-C}_5\text{H}_5)_2$  (d)  $X = \text{Ni}(\text{CO})_6$ ;  $Y = \text{Fe}(\eta^1\text{-C}_5\text{H}_5)_2$
26. The **correct** order of crystal field strength is: (Given: en = ethylenediamine)  
 (a)  $\text{Cl}^- < \text{H}_2\text{O} < \text{en} < (\eta^5\text{-C}_5\text{H}_5)^-$   
 (b)  $\text{H}_2\text{O} < \text{Cl}^- < (\eta^5\text{-C}_5\text{H}_5)^- < \text{en}$   
 (c)  $\text{H}_2\text{O} < (\eta^5\text{-C}_5\text{H}_5)^- < \text{en} < \text{Cl}^-$   
 (d)  $\text{en} < \text{Cl}^- < \text{H}_2\text{O} < (\eta^5\text{-C}_5\text{H}_5)^-$
27. The **carbon-oxygen bond** in an organic compound absorbs electromagnetic radiation of frequency  $6 \times 10^{13}$  Hz. This frequency corresponds to the region:  
 (a) Infrared (b) Microwave (c) Ultraviolet (d) Visible
28. According to the equipartition principle of energy, the molar heat capacity at constant volume for  $\text{CO}_2(\text{g})$ ,  $\text{SO}_2(\text{g})$  and  $\text{H}_2\text{O}(\text{g})$  follows the trend:  
 (a)  $\text{CO}_2 = \text{SO}_2 = \text{H}_2\text{O}$  (b)  $\text{CO}_2 > \text{SO}_2 = \text{H}_2\text{O}$   
 (c)  $\text{H}_2\text{O} > \text{SO}_2 = \text{CO}_2$  (d)  $\text{CO}_2 = \text{SO}_2 > \text{H}_2\text{O}$
29.  $\left[ \frac{-h^2}{(8\pi^2m)} \frac{d^2}{dx^2} + \frac{h^2\alpha^2x^2}{(2\pi^2m)} \right] \exp(-\alpha x^2) = C \frac{h^2}{(4\pi^2)} \exp(-\alpha x^2)$ , where  $h, \pi, m$  and  $\alpha$  are constants. Then C is:  
 (a)  $\frac{2\alpha}{m}$  (b)  $\frac{\alpha}{2m}$  (c)  $\frac{\alpha}{m}$  (d)  $\frac{\alpha^2}{m}$
30. Among Ar,  $\text{NH}_4\text{Cl}$ , HF and HCl, the strength of interatomic / intermolecular forces follows the order:  
 (a)  $\text{NH}_4\text{Cl} > \text{HF} > \text{HCl} > \text{Ar}$   
 (b)  $\text{HF} > \text{HCl} > \text{Ar} > \text{NH}_4\text{Cl}$   
 (c)  $\text{HCl} > \text{Ar} > \text{NH}_4\text{Cl} > \text{HF}$   
 (d)  $\text{Ar} > \text{NH}_4\text{Cl} > \text{HF} > \text{HCl}$
31. The number of **degrees of freedom** in the homogeneous liquid region of a two component system with a **eutectic point**, at one atmosphere pressure, is:  
 (a) 0 (b) 1 (c) 2 (d) 3
32. The **ionic strength** of 0.1 M aqueous solution of  $\text{Fe}_2(\text{SO}_4)_3$  is:  
 (a) 0.1 M (b) 0.65 M (c) 1.3 M (d) 1.5 M
33. If the transport number of  $\text{Na}^+$  is 0.463 (dilute solution of NaCl in MeOH), the transport number of  $\text{H}^+$  (dilute solution of HCl in MeOH) is: ( $\Lambda^\infty$  (NaCl in MeOH) =  $96.9 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  and  $\Lambda^\infty$  (HCl in MeOH) =  $192 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ )  
 (a) 0.27 (b) 0.46 (c) 0.54 (d) 0.73



34. Charcoal (**1 gram**) of surface area  $100 \text{ m}^2$  per gram, absorbs 60mg of acetic acid from an aqueous solution at  $25^\circ\text{C}$  and 1 atmosphere pressure. The number of moles of acetic acid adsorbed per  $\text{cm}^2$  of charcoal surface is:
- (a)  $10^{-2}$  (b)  $10^{-6}$  (c)  $10^{-5}$  (d)  $10^{-9}$
35. The change in entropy for the following transformations is respectively:  
(+ indicates increase, – indicates decrease and 0 indicates no change)



(iv) Adiabatic reversible expansion of an ideal gas

- (a) +, –, 0, + (b) +, –, 0, 0 (c) –, +, +, 0 (d) +, –, +, 0

**Attempt ALL the questions. Questions 36 – 43 (subjective questions) carry fifteen marks each.**

36. Using crystal field theory (CFT), for the  $[\text{Co}(\text{NH}_3)_6]^{3+}$  ion
- (a) draw the d-orbital splitting including their orbital labels (designations) and show their electron occupancy.
- (b) calculate the crystal field stabilization energy (ignore pairing energy) and spin-only magnetic moment values. (Given: atomic number of Co = 27).
37. (a) Write the correct order of lattice energy for LiX, X = F, Cl, Br and I.
- (b) A first order reflection from (111) plane is observed for LiX with  $2\theta = 24.6^\circ$  (X-ray of wavelength  $1.54 \text{ \AA}$ ). Assuming LiX to be a cubic crystal system, calculate the length of the side of the unit cell in  $\text{\AA}$ .
38. For the reaction:




- (i) Write the expression for the rate of the reaction in terms of the change in concentrations of NO and  $\text{H}_2\text{O}$ .
- (ii) Given the following data for the above reaction, find the order of the reaction with respect to (a) NO and (b)  $\text{H}_2$  and the rate constant of the reaction along with the proper unit.



	$[\text{NO}]_{t=0}(\text{mol dm}^{-3})$	$[\text{H}_2]_{t=0}(\text{mol dm}^{-3})$	Initial rate ( $\text{mol dm}^{-3} \text{s}^{-1}$ )
Experiment 1	0.025	0.01	$2.4 \times 10^{-6}$
Experiment 2	0.025	0.005	$1.2 \times 10^{-6}$
Experiment 3	0.0125	0.01	$0.6 \times 10^{-6}$

39. The vapour pressure of benzene is 5333 Pa at 7.6 °C and 53330 Pa at 60.6 °C. Calculate the heat of vaporization of benzene and the normal boiling point of benzene.
40. The following graph represents the dependence of certain properties I to V (given below) as a function of temperature.



Property

- I. The enthalpy change of a gas phase reaction in which the sum of the number of moles of products is greater than the sum of the number of moles of reactants
- II. The osmotic pressure of an ideal solution at a given concentration
- III. The standard Gibbs free energy of formation of metal oxides
- IV. The molar heat capacity at constant volume for an ideal gas, as predicted by the equipartition of energy
- V. The rate constant of a reaction with  $E_a = 100 \text{ kJ mol}^{-1}$

The lines / curves A, B, C, D and E corresponding to the appropriate property are:

Answer:


property	line / curves
I	
II	






III	
IV	
V	

41. Draw the structures A-E for the given transformation:




42. In the reaction sequence given below, draw the structures of A, C, D and reagent B.



43. (a) How many  $^1\text{H-NMR}$  signals are expected for 2-chlorobut-2-ene? (ignore spin-spin coupling)

(b) Write down the iron containing chemical species, E, F and G in the following reactions.



### Answer Key

Q.No	Ans		Q.No	Ans		Q.No	Ans		Q.No	Ans
1.	a		11.	b		21.	d		31.	c
2.	b		12.	c		22.	b		32.	d
3.	c		13.	d		23.	c		33.	d
4.	d		14.	b		24.	c		34.	d
5.	c		15.	d		25.	a		35.	d
6.	a		16.	a		26.	a			
7.	d		17.	c		27.	a			

## JAM – 2014 – CY

8.	c		18.	a		28.	b			
9.	d		19.	b		29.	c			
10.	b		20.	a		30.	a			

- © No Part of this Question Paper shall be reproduced, reprinted or Translated for any purpose whatsoever without prior permission of AJ Chem Academy.
- © In spite of best efforts taken to present this Work without mistakes, some mistakes may have inadvertently crept in. So, we do not take any legal responsibility for them. If they are brought to our notice, corrections will be done in next edition.

- © இந்த வினாத்தாளின் எந்தப் பகுதியும் ஏஜே கெம் அகாடமியின் முன் அனுமதியின்றி எந்த நோக்கத்திற்காகவும் மீண்டும் உருவாக்கப்படவோ, மறுபதிப்பு செய்யவோ அல்லது மொழிபெயர்க்கவோ கூடாது.
- © இந்த படைப்பை பிழையின்றி வழங்குவதற்கு சிறந்த முயற்சிகள் எடுக்கப்பட்டாலும், சில தவறுகள் கவனக்குறைவாக ஊடுருவியிருக்கலாம். எனவே அவற்றிற்கு நாங்கள் எந்த சட்டப் பொறுப்பையும் ஏற்கவில்லை. அவற்றை எங்கள் கவனத்திற்கு கொண்டு வந்தால், அடுத்த பதிப்பில் திருத்தங்கள் செய்யப்படும்.

